

Table SII. Comparison of laboratory data between patients with and without reactive skin lesions in patients with adult-onset immunodeficiency due to anti-interferon- γ autoantibodies

		<i>n</i>	Mean \pm SD	<i>p</i>
Hemoglobin (g/dl)	Without RS	141	11.29 \pm 1.94	0.751
	With RS	23	10.04 \pm 2.07	
White blood cell count (*10 ³ /ul)	Without RS	140	12.47 \pm 7.67	<0.001
	With RS	23	25.09 \pm 13.67	
Absolute neutrophil count (*10 ³ /ul)	Without RS	141	8.65 \pm 7.06	<0.001
	With RS	23	20.03 \pm 12.55	
Absolute lymphocyte count (*10 ³ /ul)	Without RS	141	2.62 \pm 1.25	0.730
	With RS	23	2.97 \pm 1.23	
Absolute eosinophil count (*10 ³ /ul)	Without RS	141	0.64 \pm 0.69	0.004
	With RS	23	1.01 \pm 1.04	
Platelet count (*10 ³ /ul)	Without RS	186	349.04 \pm 150.07	0.989
	With RS	34	431.94 \pm 134.20	

Statistical differences between the two groups were analyzed by paired *t*-test and was completed by GraphPad Prism 6 software (GraphPad Software Inc., La Jolla, CA).
RS: reactive skin diseases.