

**Table SI. Main clinical features of 20 patients with Becker's naevus**

No.	Age/duration, years	Sex	Location	Hairiness
1	15/10	Male	Right arm	Yes
2 <sup>a</sup>	19/5	Male	Left jaw and neck	No
3	16/1	Male	Left thigh and buttock	Yes
4	14/1	Male	Left upper back	No
5 <sup>a</sup>	21/3	Male	Right neck	Yes
6 <sup>a</sup>	15/14	Male	Left and right abdomen	Yes
7 <sup>a</sup>	19/6	Male	Right shoulder and chest	Yes
8 <sup>a</sup>	17/5	Male	Left chest	Yes
9 <sup>a</sup>	17/5	Male	Left waist and buttock	Yes
10 <sup>a</sup>	16/12	Male	Right thigh	Yes
11 <sup>a</sup>	17/7	Male	Left and right waist	Yes
12 <sup>a</sup>	16/4	Male	Left scapular region	Yes
13 <sup>a</sup>	12/6	Male	Left scapular region	Yes
14	13/0.1	Male	Left forearm and right thigh	No
15	28/14	Male	Right scapular region and arm	Yes
16 <sup>a</sup>	23/5	Male	Right thigh	Yes
17 <sup>a</sup>	38/32	Male	Left thigh	Yes
18 <sup>a</sup>	25/10	Female	Right scapular region	Yes
19	23/20	Female	Right scapular region	Yes
20	17/10	Female	Right thigh	Yes

<sup>a</sup>Reverse transcription PCR (RT-PCR) and Western blot were performed on lesional and perilesional skin from these 13 cases.