

Table SIII. Immunostaining scores of androgen receptor (AR), oestrogen receptor (ER) α , ER β and progesterone receptor (PR) in Becker's naevus and normal controls

		Interfollicular epidermis				Hair follicles				Sebaceous glands			
		<i>n</i>	1+	2+	3+	<i>n</i>	1+	2+	3+	<i>n</i>	1+	2+	3+
AR	Lesion	20	1	17	2	14	2	12	0	12	0	8	4
	Perilesion	20	5	14	1	11	9	2	0**	11	1	10	0*
	Control	10	5	5	0**								
ER α	Lesion	20	0	6	14	14	0	3	11	12	0	5	7
	Perilesion	20	1	18	1**	11	0	10	1**	11	0	6	5
	Control	10	4	6	0** Δ								
ER β	Lesion	20	0	6	14	14	0	4	10	12	0	8	4
	Perilesion	20	1	14	5**	11	0	8	3*	11	0	9	2
	Control	10	3	7	0** Δ								
PR	Lesion	20	0	1	19	14	0	1	13	12	0	3	9
	Perilesion	20	0	15	5**	11	3	5	3**	11	0	3	8
	Control	10	3	7	0** $\Delta\Delta$								

* $p < 0.05$, ** $p < 0.01$, compared with lesional skin; $\Delta p < 0.05$, $\Delta\Delta p < 0.01$, compared with perilesional skin.