

Table S1. Clinicopathological features of patients with angioimmunoblastic T-cell lymphoma (AITL) who developed secondary Epstein-Barr virus (EBV)-associated diffuse large B-cell lymphoma (DLBCL)

Case No.	Authors	Age, years/sex	Interval, months	Tumour site		EBV status in AITL	Treatment		Clinical outcome
				AITL	DLBCL		AITL	DLBCL	
1	Abruzzo et al. (1993) (S1)	46/M	26	LN	Skin, LN and spleen	-	Cyclophosphamide	N/D	Died of multi-organ failure
2	Matsue et al. (1998) (S2)	55/M	35	LN	LN	+	CC	CEPP	Died 13 months later of multi-organ failure
3		58/M	20	LN	Lung, spleen, BM, LN, liver, and submandibular gland	-	CHOP+BMT	N/D	Died 8 months later of multi-organ failure
4	Park et al. (2002) (9)	55/F	48	LN	LN	N/Av	CHOP	CHOP-R	CR and alive 12 months later
5	Zetti et al. (2002) (8)	68/M	34	LN	Soft tissue	+	Observation	Glucocorticoids	Died 4 months later of tuberculosis
6		47/M	29	LN	LN	+	CC+BMT	CC+RT	Died 3 months later of Aspergillus pneumonia
7	Hawley et al. (2006) (S3)	69/M	56	LN	Skin	-	CHOP	R-hyperCVAD	Died 54 months later
8	Attygalle et al. (2007) (S4)	28/M	24	LN	BM	N/Av	CHOP	Gemcitabine	Died 3 months later
9		60/F	8	LN	LN	+	CHOP	CHOP-ET	Died 3 months later
10		59/M	8	LN	LN	N/Av	N/Av	CC	PR and alive with disease 68 months later
11	Willenbrock et al. (2007) (S5)	72/M	84	LN	LN	N/Av	N/Av	N/Av	N/Av
12		76/F	8	LN	LN	N/Av	N/Av	N/Av	N/Av
13		52/F	3	LN	LN	N/Av	N/Av	N/Av	N/Av
14		46/F	5	LN	BM	N/Av	N/Av	N/Av	N/Av
15	Weisel et al. (2008) (12)	59/M	11	LN	DB and lung	-	Fudarabine+CHOP	N/D	Died 2 weeks later
16	Takahashi et al. (2010) (10)	66/F	24	LN	Ileum	+	THPCOP+CHASE	N/D	Died 1 months later of respiratory failure
17	Skugor et al. (2010) (S6)	36/F	11	LN	LN	N/Av	FED	CHOP-R+BMT	CR
18	Huang et al. (2011) (S7)	64/M	47	LN	LN	+	IHOP	CHOP-R	Alive with disease 13 months later
19	Yang et al. (2012) (7)	65/M	19	LN	Skin	+	CHOP	CHOP-R	PR
20	Smeltzer et al. (2012) (11)	59/F	18	LN	Bilateral AG	+	CHOP+BMT+Vorinostat	R	CR
21	Present case	49/F	22	LN	Skin	-	CHOP	CHOP-R	Died 5 months later of multi-organ failure

LN: lymph node; BM: bone marrow; DB: duodenal bulb; AG: adrenal gland; N/Av: not available; N/D: not done; CC: combination chemotherapy; details not available; CEPP: endoxan; etoposide; procarbazine and prednisolone; CHOP: cyclophosphamide; vincristine; adriamycin; and prednisone; BMT: bone marrow transplantation; R: rituximab; RT: radiation treatment; hyperCVAD: cyclophosphamide; vincristine; doxorubicin; and dexamethasone; Et: etoposide; THPCOP: cyclophosphamide; pirarubicin; vincristine; prednisolone; CHASE: cyclophosphamide; cytarabine, etoposide; dexamethasone; FED: fludarabine; cyclophosphamide; dexamethasone; IHOP: ifosfamide; doxorubicin; vincristine; and prednisone; ESHAP: etoposide; methyl prednisolone; cytarabine (ara-C) and cisplatin; PR: partial response; CR: complete remission.

SUPPLEMENTARY REFERENCES

1. Abruzzo LV, Schmidt K, Weiss LM, et al. B-cell lymphoma after angioimmunoblastic lymphadenopathy: a case with oligo clonal gene rearrangements associated with Epstein-Barr virus. *Blood* 1993; 82: 241-6.
2. Matsue K, Itoh M, Tsukuda K, Kokubo T, Hirose Y. Development of Epstein-Barr virus-associated B cell lymphoma after intensive treatment of patients with angioimmunoblastic lymphadenopathy with dysproteinemia. *Int J Hematol* 1998; 67: 319-29.
3. Hawley RC, Cankovic M, Zarbo RJ. Angioimmunoblastic T-cell lymphoma with supervening Epstein-Barr virus-associated large B-cell lymphoma. *Arch Pathol Lab Med* 2006; 130: 1707-11.
4. Attygalle AD, Kyriakou C, Dupuis J, et al. Histologic evolution of angioimmunoblastic T-cell lymphoma in consecutive biopsies: clinical correlation and insights into natural history and disease progression. *Am J Surg Pathol* 2007; 31: 1077-88.
5. Willenbrock K, Bräuninger A, Hansmann ML. Frequent occurrence of B-cell lymphomas in angioimmunoblastic T-cell lymphoma and proliferation of Epstein-Barr virus-infected cells in early cases. *Br J Haematol* 2007; 138: 733-9.
6. Skugor ND, Perić Z, Vrhovac R, Radić-Kristo D, Kardum-Skelin I, Jaksčić B. Diffuse large B-cell lymphoma in patient after treatment of angioimmunoblastic T-cell lymphoma. *Coll Antropol* 2010; 34: 241-5.
7. Huang J, Zhang PH, Gao YH, Qiu LG. Sequential development of diffuse large B-cell lymphoma in a patient with angioimmunoblastic T-cell lymphoma. *Diagn Cytopathol* 2012; 40: 346-51.