

Table SII. Clinical signs and symptoms in the full cohort of 844 patients with papulopustular rosacea (PPR)-suggestive or demodicosis, and in the two subgroups^a

Clinical signs and symptoms	Full cohort n = 844 n (%)	Subgroup A n = 488 n (%)	Subgroup B n = 490 n (%)
Vascular symptoms ^b	677 (80)	366 (75)	399 (81)
Flushes	531 (63)	287 (59)	316 (64)
Persistent erythema	601 (71)	325 (67)	353 (729)
Follicular scales (skin)	782 (93)	457 (94)	460 (94)
Papules	330 (39)	177 (36)	174 (36)
Pustules	188 (22)	101 (21)	91 (19)
Follicular eczematids	108 (13)	41 (8)	75 (15)
Pigmentation	15 (2)	8 (2)	12 (2)
Cylindrical dandruff (lash)	72 (9)	65 (13)	76 (16)
Blepharitis	78 (9)	56 (12)	56 (11)
Conjunctivitis	12 (1)	7 (1)	9 (2)
Chalazion	18 (2)	12 (3)	9 (2)
Pruritus	130 (15)	117 (24)	68 (14)
Scalp complaint	321 (38)	244 (50)	164 (33)
Ocular involvement	180 (21)	140 (29)	103 (21)

^aA (488 patients included from August 2005) and B (490 patients who had had no recent treatment and did not have an associated facial dermatosis). ^bflushes or persistent erythema.