

Table SII. Differences in profession, years of clinical experience and practice setting for respondents prescribing antiepileptics or antidepressants for chronic pruritus

	Respondents prescribing antiepileptic treatment (<i>n</i> = 73) <i>n</i> (%)	Respondents not prescribing antiepileptic treatment (<i>n</i> = 119) <i>n</i> (%)	<i>p</i> -value
Prescribing antiepileptics			
Profession			0.002
Dermatology resident	23 (60.5)	15 (39.5)	
Dermatologist	51 (32.9)	104 (67.1)	
Years of clinical experience			0.001
0–10 years	49 (51.0)	49 (49.0)	
10–20 years	12 (24.0)	38 (76.0)	
> 20 years	12 (27.7)	34 (72.3)	
Practice setting			0.000
Academic hospital (<i>n</i> = 50)	33 (66.0)	17 (34.0)	
General hospital (<i>n</i> = 101)	27 (26.7)	74 (73.3)	
Private practice (<i>n</i> = 21)	4 (19.0)	17 (81.0)	
Other (<i>n</i> = 4)	1 (25.0)	3 (75.0)	
More than one practice setting (<i>n</i> = 17)	9 (52.9)	8 (47.1)	
Prescribing antidepressants			
	Respondents prescribing antidepressants (<i>n</i> = 101)	Respondents not prescribing antidepressants (<i>n</i> = 91)	
Profession			0.466
Dermatology resident	22 (57.9)	16 (42.1)	
Dermatologist	79 (51.3)	75 (48.7)	
Years of clinical experience			0.346
0–10 years (<i>n</i> = 95)	55 (57.9)	40 (42.1)	
10–20 years (<i>n</i> = 50)	24 (48.0)	26 (52.0)	
> 20 years (<i>n</i> = 47)	22 (46.8)	25 (53.2)	
Practice setting			0.241
Academic hospital (<i>n</i> = 50)	33 (66.0)	17 (34.0)	
General hospital (<i>n</i> = 101)	50 (49.5)	51 (50.5)	
Private practice (<i>n</i> = 21)	8 (40.0)	12 (60.0)	
Other (<i>n</i> = 4)	2 (50.0)	2 (50.0)	
More than one practice setting (<i>n</i> = 17)	8 (47.1)	9 (52.9)	