

Table I. Summary of clinical and histological characteristics of the vulvar melanomas among patients with or without lichen sclerosus (LS)

Case no.	Age, LS years	Relapse, years after diagnose	Site	Tumour size, mm	Tumour depth, mm	Clark T N	M Stage	Ulceration	Lymph node status	Treatment	Survival
1	+	69	Left labium	30	19	(V)	4b X 1 IV	+	NS	Vulvectomy and reconstructive surgery	DOD
2	+	59	Right minor labium	40	10	IV	4b 2b 1 IV	+	3/12	Wide dissection, node resection	DOD
3	+	79	Right major labium	NS (size of finger-tip)	9	IV	4b X IV	+	NS	Local resection	DOD
4	-	67	Labium minus left	8	9	III	4b 0 IIIC	+	0/2	Local resection	DOD
5	-	80	Local metastasis	NS	NS	NS					
6	-	83	Right minor labium	20	11	IV	4b 0 0 IIIC	+	0/3	Local resection	DOD
			Hymenal ring, vaginal collarette	NS	NS	NS				Brachytherapy at residive	
			Left labium	NS	1.6	IV	2b 2b 0 III	+	2/3	Local resection	
7	-	66	Clitoris	13	1.2	III	2a 0 0 IB	-	0/3	Local resection	DOD
8	-	64	Anterior to urethra	40	NS	NS			6/18		
			Left labium	15	2	IV	3b 0 0 IIIB	+	0/2	Local resection	AWOD
9	-	75	Midline	9	9	V	4b 0 IIIC	+	0/1	Local resection	AWOD

Lymph node status: number of positive lymph nodes/number of lymph nodes studied.

NS: not studied; DOD: died of disease; AWOD: alive without disease.