

Table SIV. Characteristics of the 12 chronic spontaneous urticaria (CSU) patients with serum samples collected in the remission phase

Age, years, median (IQR)	33 (30; 41)
Sex (female/male)	9/3
Female age, years, median (IQR)	35 (31.5; 45)
Male age, years, median (IQR)	30 (27; 33)
Disease duration, months, median (IQR)	3.5 (2.3; 5.0)
Personal history of atopic diseases, <i>n</i> (%)	2 (16.7)
Urticaria/atopic diseases family history, <i>n</i> (%)	8 (66.7)
Urticaria family history, <i>n</i>	4
Atopic diseases family history, <i>n</i>	4
Urticaria and atopic diseases family history, <i>n</i>	0
UAS, median; (IQR)	5 (4; 6)
Total IgE, IU/ml, median (IQR)	142.7 (39.6; 365.7)
MiR-125a-5p ($2^{-\Delta\Delta Ct}$), median (IQR)	
Active phase	0.028 (0.007; 0.058)
Remission phase	0.004 (0.002; 0.009)
CCL17, pg/ml, median (IQR)	
Active phase	597.1 (346.3; 976.6)
Remission phase	343.3 (256.6; 488.9)
Positive anti-TG/anti-TPO antibodies, <i>n</i> (%)	2 (16.7)
Positive serum specific IgE, <i>n</i> (%)	4 (33.3)

HC: healthy control; IQR: interquartile ranges; UAS: urticaria activity score; TG: thyroglobulin; TPO: thyroid peroxidase.