

Table S1. Reported cases of neutrophilic dermatoses (ND) induced by pegfilgrastim

Case	Authors	Age, years/ Sex	Type of ND	Primary disease	Past usage of filgrastim	Initial use of pegfilgrastim	Time between administration and onset	Location of symptom	Fever	WBC (x10 ³ /μl)	Treatment	Time to cure	Recurrence
Case 1	Draper et al. (6)	48/M	Bullous SS	Severe congenital neutropenia	Yes	Yes	3 days	Face, trunk, extremities	Low-grade fever	24.7	Prednisone 60 mg/day	2 weeks	None
Case 2	Llamas-Velasco et al. (7)	77/F	SS with subcutaneous involvement	Acute myeloid leukemia	No	Yes	10 days	Back, both arms, left leg	Fever	0.12	Prednisone 40 mg/day	1 week	None
Case 3	Machan et al. (8)	43/F	SS	Drug-induced neutropenia (Ziprasidone)	No	Yes	10 days	Trunk and extremities	Fever	5.5	Intravenous methylprednisolone	2 days	None
Case 4	White et al. (9)	23/M	Pyoderma gangrenosum	Hodgkin's disease	Yes	Yes	3 days	Right upper arm, chest, shoulder, forehead	Fever	15.3	IVIG, prednisone 1 mg/kg/day	3 weeks	None
Case 5 (present case)	Present study	59/F	ND comparable to SS	Chemotherapy-induced neutropenia (breast cancer)	Yes	Yes	10 days	Right ankle	None (37.1°C)	18	Cessation of pegfilgrastim	2 weeks	None

SS: Sweet's syndrome; WBC: white blood cells