

APPENDIX S1

ACKNOWLEDGEMENTS

BEST Study Group: Dr Per Åkerlund, Falu Lasarett, Falun, Sweden; Dr Tun Aung, West Park Rehabilitation Hospital, Wolverhampton, UK; Dr Janne Biörklund, Rehabcentrum Lund-Orup, Lund, Sweden; Professor Jörgen Borg, Department of Clinical Sciences, Karolinska Institutet, Rehabilitation Medicine, Danderyd Hospital, Stockholm, Sweden; Professor Dr Andres Ceballos-Baumann, Neurologisches Krankenhaus München, München, Germany; Dr Christian Dohle, MED IAN Klinik Berlin, Abteilung für Neurologische Rehabilitation, Charité – Universitätsmedizin, Berlin, Germany; Dr Micael Edbom, Länssjukhuset Ryhov, Jönköping, Sweden; Dr Per Ertzgaard, Department of Rehabilitation Medicine, Linköping University, Linköping, Sweden; Dr Naseer Haboubi, Linden Lodge Rehabilitation Centre, Nottingham, UK; Dr Anders Häggström, Rehabiliteringsmedicinska Kliniken, Örebro, Sweden; Dr Christoph Herrmann, Clinics for Neurological Rehabilitation and Early Rehabilitation, Asklepios Clinics Schildautal, Seesen, Germany; Professor Stefan Hesse, Medical Park Berlin Humboldtmühle Neurologie, Berlin, Germany; Professor Jai Kulkarni, Manchester Royal Infirmary, Manchester, UK; Dr Kjell Kullander, Karolinska Universitetssjukhuset, Huddinge, Sweden; Dr Lal Landham, Medway Maritime Hospital, Gillingham, UK; Dr Kristina Lindgren, Centralsjukhuset, Neuro-och rehabkliniken, Karlstad, Sweden; Dr Sabine Mehnert, Justus-Liebig University, Giessen, Germany; Dr Friedmann Müller, Neurologische Klinik Bad Aibling, Bad Aibling, Germany; Dr Margaret Phillips,

Derby City Hospital, Derby, UK; Professor Thomas Platz, BDH-Klinik und Ernst-Moritz-Arndt Universität Greifswald, Germany; Dr Robert Prempeh, Falkirk & District Royal Infirmary, Falkirk, UK; Dr Iris Reuter, Justus-Liebig University, Giessen, Germany; Dr Chris Roy, Southern General Hospital, Glasgow, UK; Dr Mohamed Sakel, Kent & Canterbury Hospital, Canterbury, UK; Dr Patrik Säterö, SU/Högsbo sjukhus, Göteborg, Sweden; Dr Lalith Satkunam, Glenrose Rehabilitation Hospital, Edmonton, Canada; Dr Wilfried Schupp, Fachklinik Herzogenaurach, Herzogenaurach, Germany; Dr Klaus Seigel, Akademiska sjukhuset, Uppsala, Sweden; Dr Satyendra Sharma, Sunnybrook Health Science Centre, Toronto, Canada; Dr Christian van der Ven, Neurologisches Rehabilitationszentrum Bonn Godeshöhe, Bonn, Germany; Dr Martina Walsh, West Midlands Rehabilitation Centre, Birmingham, UK; Professor Anthony B. Ward, North Staffordshire Rehabilitation Centre, University Hospital of North Staffordshire, Stoke on Trent, UK; Dr Theodore Wein, Montreal General Hospital, Montreal, Canada; Professor Dr Jörg Wissel, Neurological Rehabilitation, Department of Neurology, Vivantes Klinikum Spandau, Berlin, Germany.

Funding: Allergan employees were involved in: (i) the design and conduct of the study; and (ii) the collection, management, analysis, and interpretation of the study data. The authors had sole control over the preparation, review, and approval of this manuscript. Editorial assistance for preparation of this manuscript was provided by Right Angle Communications, funded by Allergan.