

Supplementary material to article by G. Francisco et al. “Adult spasticity international registry study: methodology and baseline patient, healthcare provider, and caregiver characteristics”

Table SII. Spasticity treatment modalities at baseline by upper limb presentation

	Adducted internally rotated shoulder (n = 185)	Clenched fist (n = 341)	Flexed elbow (n = 348)	Flexed wrist (n = 267)	Intrinsic plus hand (n = 81)	Pronated forearm (n = 163)	Thumb-in-palm (n = 109)	Other (n = 49)	Total (n = 527)
Acupuncture, n (%)	183 16 (8.7)	330 31 (9.4)	335 28 (8.4)	259 16 (6.2)	78 7 (9.0)	160 12 (7.5)	107 14 (13.1)	46 6 (13.0)	505 43 (8.5)
Assistive Devices, n (%)	183 144 (78.7)	330 248 (75.2)	335 242 (72.2)	259 189 (73.0)	78 55 (70.5)	160 115 (71.9)	107 70 (65.4)	46 36 (78.3)	505 367 (72.7)
Casting, n (%)	183 13 (7.1)	330 31 (9.4)	335 36 (10.7)	259 30 (11.6)	78 12 (15.4)	160 18 (11.3)	107 8 (7.5)	46 7 (15.2)	505 56 (11.1)
Chemodenervation, n (%)	183 15 (8.2)	330 21 (6.4)	335 17 (5.1)	259 8 (3.1)	78 10 (12.8)	160 11 (6.9)	107 6 (5.6)	46 2 (4.3)	505 25 (5.0)
Intrathecal therapy, n (%)	183 25 (13.7)	330 39 (11.8)	335 34 (10.1)	259 27 (10.4)	78 13 (16.7)	160 23 (14.4)	107 13 (12.1)	46 8 (17.4)	505 60 (11.9)
Orthotics, n (%)	183 120 (65.6)	330 209 (63.3)	335 210 (62.7)	259 156 (60.2)	78 55 (70.5)	160 112 (70.0)	107 63 (58.9)	46 26 (56.5)	505 298 (59.0)
Physio or occupational therapy, n (%)	183 158 (86.3)	330 284 (86.1)	335 277 (82.7)	259 220 (84.9)	78 68 (87.2)	160 142 (88.8)	107 89 (83.2)	46 40 (87.0)	505 420 (83.2)
Surgeries or other procedures, n (%)	183 22 (12.0)	330 35 (10.6)	335 35 (10.4)	259 24 (9.3)	78 11 (14.1)	160 15 (9.4)	107 12 (11.2)	46 7 (15.2)	505 64 (12.7)

NOTE: Allowed responses included “yes” or “not mentioned”.