

Table SIII. Attrition analysis descriptive summary

| Variable | Completers n = 51 | Non-completers n = 14 | Statistic | p-value |
|------------------------------|----------------------|--------------------------|-----------|---------|
| Participant variables | | | | |
| Age (years) | 61.94 (11.80) | 56.92 (15.73) | t = 1.30 | 0.19 |
| % Female | 43.14 | 35.71 | χ^2 | 0.62 |
| Education (years) | 14.22 (2.37) | 14.00 (2.69) | t = 0.47 | 0.77 |
| Estimated IQ | 103.02 (12.41) | 101.23 (11.33) | t = -0.25 | 0.64 |
| MoCA total | 24.22 (2.87) | 24.07 (2.50) | t = 0.18 | 0.86 |
| NEADL total | 19.60 (2.95) | 20.40 (2.07) | t = -0.82 | 0.42 |
| Stroke variables | | | | |
| Time since stroke | 41.04 (43.01) | 44.75 (53.81) | t = 0.29 | 0.80 |
| % Left hemisphere | 58.00 | 41.67 | χ^2 | 0.50 |
| % Ischaemic | 66.67 | 78.57 | χ^2 | 0.49 |
| Objective memory | | | | |
| Verbal learning ^a | -0.65 (1.34) | -0.61 (1.36) | t = -0.09 | 0.93 |
| Verbal recall ^a | -0.84 (1.43) | -0.88 (1.36) | t = 0.08 | 0.93 |
| Visual learning ^a | -0.77 (1.20) | -1.16 (1.53) | t = 1.00 | 0.31 |
| Visual recall ^a | -0.64 (1.29) | -1.24 (1.96) | t = 1.35 | 0.18 |
| Verbal WM ^b | 10.82 (3.51) | 9.93 (2.27) | t = 0.90 | 0.37 |
| Visual WM ^b | 8.39 (2.21) | 9.00 (2.00) | t = -0.93 | 0.36 |
| Strategy use | | | | |
| External | 12.12 (5.82) | 13.83 (5.92) | t = -0.91 | 0.36 |
| Internal | 3.96 (3.53) | 3.17 (3.51) | t = 0.70 | 0.49 |
| Subjective memory | | | | |
| Everyday memory | 21.02 (13.54) | 12.88 (9.17) | t = 1.40 | 0.11 |
| PM-self | 65.58 (22.25) | 49.89 (11.93) | t = 2.05 | 0.05 |
| PM close other | 60.79 (23.69) | 66.00 (13.21) | t = 1.63 | 0.55 |

^aValues are z-scores.

^bValues are aged scaled [AQ5] scores.

IQ: intelligence quotient; MoCA: Montreal Cognitive Assessment; NEADL: Nottingham Extended Activities of Daily Living; PM: prospective memory; WM: working memory; χ^2 : chi-squared test for independence of fit.