

Supplementary material to article by E. A. de Vries et al. "Fatigue in relation to long-term participation outcome in aneurysmal subarachnoid haemorrhage survivors"

Table SI. Overview of the questionnaires

Questionnaire and construct	Number of items and example(s)	Scoring	Outcomes, ranges and interpretation
FSS: Severity of fatigue	9 statements about fatigue: <i>Fatigue interferes with my physical functioning</i>	1–7: Strongly disagree – Strongly agree	Mean score 1–7, cut-off: ≥ 4 fatigued Higher score indicates more severe fatigue
SIP-68: Functional Performance in activities of daily living	68 items about functional performance: <i>I walk more slowly;</i> <i>I am doing fewer community activities</i>	One point if the item applies to the current health status of the patient	Sum score for all scales: Total scale 0–69 Physical scale 0–39 Psychosocial scale 0–29 Higher scores indicate worse functional performance
IPA: Participation autonomy, participation restrictions and problems with the level of participation	32 items, divided over 5 subscales: <i>My chances of visiting relatives and friends when I want to are...;</i> <i>My chances of spending leisure time the way I want to are...</i> 9 items about problem experience: <i>With regard to your leisure time, to what extent does your health or disability cause problems?</i>	Items of 5 subscales: 0–4: Very good – Very poor Problem experience items: 0–2: No problems – Severe problems	Mean score for each subscale: Autonomy indoors 0–4 Family role 0–4 Autonomy outdoors 0–4 Social relations 0–4 Work and Education 0–4 Problem experience 0–2 Higher scores indicate less autonomy, more restrictions and more problems
CIQ: Community Integration	13 items, divided over 3 subscales <i>Who usually prepares meals in your household?;</i> <i>How often do you travel outside the home?</i>	Home competency and Social integration items: 0–2: Alone – Together/Low – High frequency Productive activity items: 0–5: Combined score	Sum scores: Total score 0–25 Home competency (5 items) 0–10 Social integration (5 items) 0–10 Productive activity (3 items) 0–5 Higher scores indicate worse community integration
CES-D: Depression	20 items about perceived mood: <i>During the past week... I felt sad</i>	0–3: Rarely or none of the time, less than 1 day – Most or all of the time, 5–7 days	Sum score 0–60 Cut-off: ≥ 16 depression

FSS: Fatigue Severity Scale; CES-D: Center for Epidemiological Studies Depression scale; SIP-68: Sickness Impact Profile-68; IPA: Impact on Participation and Autonomy questionnaire; CIQ: Community Integration Questionnaire.