

Supplementary material to article by L. Turner-Stokes et al. “Longitudinal goal attainment with integrated upper limb spasticity management including repeat injections of botulinum toxin A: Findings from the prospective, observational Upper Limb International Spasticity (ULIS-III) cohort study”

Table SIV. Physical therapy interventions (overall)

Parameter, n (%)	Cycle 1 (n = 953) n = 818	Cycle 2 (n = 840) n = 728	Cycle 3 (n = 744) n = 575	Cycle 4 (n = 593) n = 367	Cycle 5 (n = 377) n = 207	Cycle 6 (n = 213) n = 99	Cycle 7 (n = 105) n = 25
Splinting	214 (26.2)	198 (27.2)	172 (29.9)	115 (31.3)	57 (27.5)	33 (33.3)	12 (48.0)
Shoulder supports/slings	25 (3.1)	22 (3.0)	20 (3.5)	8 (2.2)	1 (0.5)	1 (1.0)	1 (4.0)
Serial casting	17 (2.1)	7 (1.0)	5 (0.9)	1 (0.3)	1 (0.5)	1 (1.0)	1 (4.0)
Positioning of upper limb	400 (48.9)	343 (47.1)	266 (46.3)	154 (42.0)	84 (40.6)	41 (41.4)	5 (20.0)
Passive stretch	588 (71.9)	510 (70.1)	407 (70.8)	280 (76.3)	156 (75.4)	79 (79.8)	21 (84.0)
Electrical stimulation	78 (9.5)	52 (7.1)	37 (6.4)	31 (8.4)	23 (11.1)	12 (12.1)	5 (20.0)
Strength training	229 (28.0)	194 (26.6)	155 (27.0)	103 (28.1)	38 (18.4)	15 (15.2)	4 (16.0)
Task practice	138 (16.9)	117 (16.1)	82 (14.3)	64 (17.4)	25 (12.1)	11 (11.1)	3 (12.0)
Other	36 (4.4)	30 (4.1)	21 (3.7)	9 (2.5)	4 (1.9)	4 (4.0)	1 (4.0)
Missing	n = 135	n = 113	n = 169	n = 226	n = 170	n = 114	n = 80