

Supplementary material to article by L. Turner-Stokes et al. "Longitudinal goal attainment with integrated upper limb spasticity management including repeat injections of botulinum toxin A: Findings from the prospective, observational Upper Limb International Spasticity (ULIS-III) cohort study"

**Table SV.** Rates of goal achievement by domain, by cycle

Domain, n (%)	% goal achievement								
	Cycle 1	Cycle 2	Cycle 3	Cycle 4	Cycle 5	Cycle 6	Cycle 7	Cycle 8	Cycle 9
Pain	230/349 (65.9)	215/274 (78.5)	162/221 (73.3)	133/178 (74.7)	84/117 (71.8)	43/62 (69.4)	17/26 (65.4)	2/9 (22.2)	1/2 (50.0)
Involuntary movement	157/209 (75.1)	148/187 (79.1)	132/155 (85.2)	89/107 (83.2)	52/58 (89.7)	22/25 (88.0)	8/9 (88.9)	1/2 (50.0)	-
Range of movement	172/228 (75.4)	159/194 (82.0)	148/172 (86.0)	130/156 (83.3)	68/84 (81.0)	43/52 (82.7)	21/28 (75.0)	2/2 (100.0)	1/1 (100.0)
Passive function	395/548 (72.1)	348/460 (75.7)	315/388 (81.2)	239/287 (83.3)	148/178 (83.1)	60/77 (77.9)	26/41 (63.4)	9/10 (90.0)	-
Active function	122/253 (48.2)	97/168 (57.7)	104/153 (68.0)	69/108 (63.9)	50/76 (65.8)	28/41 (68.3)	19/25 (76.0)	5/6 (83.3)	-
Mobility	21/34 (61.8)	14/19 (73.7)	10/10 (100.0)	8/9 (88.9)	3/5 (60.0)	2/5 (40.0)	3/5 (60.0)	-	-
Other	27/36 (75.0)	23/33 (69.7)	14/22 (63.6)	11/15 (73.3)	8/9 (88.9)	5/5 (100.0)	0/1 (0.0)	1/1 (100.0)	-