

Table SIII. Melanocytic naevi (MN) counts or MN density among Caucasian children

Reference	Children, <i>n</i>	Age, years	Country	MN counts (mean)
Gallagher, et al. 1990 (50)	913	6–18	Canada	8.8 (6–7 years); 18 (16–18 years)
Pope, et al. 1992 (44)	1,953	4–11	UK	23.5
Green, et al. 1995 (45)	102	11–14	Australia	116.6
Luther, et al. 199 (64)	357	1–6	Germany	9 vs. 40 (5 years later)
Autier, et al. 1998 (20)	631	6–7	Europe	6
Enta & Kwan 1998 (51)	332	5–15	Canada*	2.34
Graham, et al. 1999 (5)	447	7	UK	28
Harrison, et al. 2000 (52)	115	0–3	Australia	1 (1 st year)–6 (3 rd year)
	157	0–3	UK	0 (1 st year)–2 (3 rd year)
Azizi, et al. 2000 (24)	974	7–12	Israel	22–25 (7 years); 47–48 (12 years)
Gallagher, et al. 2000 (21)	458	6–10	Canada	41 (6–7 years); 68 (9–10 years)
Dulon, et al. 2002 (53)	11,478	5–6	Germany	11
Darlington, et al. 2002 (54)	111	12–13	Australia	130 (1990); 215 (1999)
Carli, et al. 2002 (55)	3,160	13–14	Italy	11
Wiecker, et al. 2003 (2)	1,812	3–7	Germany	3 (3 years)–19 (7 years)
Synnerstad, et al. 2004 (56)	524	8–9	Sweden	8
Valiukeviciene, et al. 2005 (35)	484	1–15	Lithuania	15
Gefeller, et al. 2007 (17)	1,898	6–7	Germany	3.09–4.7
Rodvall, et al. 2007 (36)	1,360	7	Sweden	5.6–10.4
Harrison, et al. 2008 (7)	479	1–6	Australia	9 (1 year); 12 (6 years)
Aguilera, et al. 2009 (34)	180	1–15	Spain	17.5
Pettijohn, et al. 2009 (18)	681	7	USA	20.7 (<2 mm); 66.5 (≥2 mm)
Oliveria, et al. 2009 (16)	443	11	USA	8.4 (back only)
Crane, et al. 2009 (57)	870	3–8	USA	6.2 (3 years); 31.6 (8 years)

*Hutterite children with a religious-based dress code.