

Table SII. Self-reported use of dermatological pharmaceuticals (%) among responders reporting dermatological diseases or problems. Reported severe and mild problems are pooled together. Statistical significances from the logistic regression analysis for each type of reported use of dermatologicals are given. Psoriasis, other eczema and acne were compared with hand eczema (reference group) controlling for sex and age (n=1,280)

Diagnosis	Dermatological pharmaceuticals					
	Topical steroids, on prescription % (p-value)	Other pharmaceuticals, on prescription % (p-value)	Topical steroids, OTC % (p-value)	Other pharmaceuticals, OTC % (p-value)	Natural/Herbal products % (p-value)	Any dermatological pharmaceutical % (p-value)
Included in the logistic regression analyses						
Hand eczema ^a	29.5	4.4	15.3	23.8	6.3	51.6
Psoriasis	46.6 (<0.0001)	9.4 (0.0020)	13.1 NS	17.8 NS	6.3 NS	61.3 (0.0002)
Other eczema	39.9 (<0.0001)	6.5 (0.0059)	22.3 (<0.0001)	20.8 NS	4.8 NS	62.4 (<0.0001)
Acne	10.8 (0.0014)	2.8 NS	8.2 NS	16.6 NS	4.8 NS	31.2 (0.0017)
Not included in the logistic regression analyses						
Urticaria	26.7	6.7	14.0	21.3	8.7	50.7
Other skin problems	26.6	7.2	10.0	24.1	6.9	51.9

^aComparison group.

OTC: over the counter; NS: not significant.