

Table SII. Mean values for estimated glomerular filtration rate (eGFR) and creatinine in the hospital HS group, population HS group, and population non-HS group and unadjusted eGFR levels according to CKD stages (p-value < 0.001).

	Hospital HS group n=32	Population HS group n=430	Population non-HS group n=20,780
Mean values for eGFR and creatinine			
eGFR (unadjusted), ml/min/1.73m ² , mean ± SD (range)	104.79 ± 14.39 (71.92–127.52)	93.60 ± 16.66 (10.64–149.55)	87.17 ± 16.46 (5.45–141.22)
eGFR (age-sex-smoking adjusted), ml/min/1.73m ² , mean (95%CI)	94.63 (89.99–99.27)	88.00 (86.72–89.29)	88.41 (88.10–88.71)
eGFR (age-sex-smoking-BMI-hypertension-diabetes adjusted), ml/min/1.73 m ² , mean (95% CI)	95.53 (90.66–100.40)	90.25 (88.55–91.96)	88.72 (88.10–89.34)
Creatinine, umol/l) mean (95% CI)	64.2 (60.1–68.3)	73.2 (71.1–75.3)	76.5 (76.2–76.8)
Unadjusted eGFR levels			
eGFR ≥ 90 ml/min/1.73 m ²	84.38%	63.64%	46.57%
eGFR 60–89 ml/min/1.73 m ²	15.63%	32.87%	47.52%
eGFR 30–59 ml/min/1.73 m ²	0%	3.03%	5.68%
eGFR 15–29 ml/min/1.73 m ²	0%	0.23%	0.18%
eGFR < 15 ml/min/1.73 m ²	0%	0.23%	0.05%