

Table SI. Clinical data and BP180 antibody levels for patients with pemphigoid gestationis (PG)

Patient	Gravidity	Parity	GA at onset of symptoms	BP180 <sup>a</sup> in first trimester of pregnancy with PG	Highest BP180 <sup>a</sup> during the course of PG	BP180 <sup>a</sup> in first trimester of subsequent pregnancies	Time after PG (months, from last PG delivery)	
1	3	2	36+	2.6	43			
	4	3					5.7	13
	5	3					3.5	19
2	3	2	32+	3.1	NA			
	4	3	10+	30	NA		24	
	5	4				25	21	
	6	5				19	41	
	7	6				17	60	
3	1	0	37+	1.9	NA			
	2	1				13	20	
	3	1	8+	51	83		23	
4	3	1	16+	3.7	NA			
	4	2				2.3	62	
5	1	0	37+	2.1	27			
	2	1				3.0	39	
	3	2				2.9	57	
6	2	1	24+	4.3	50	No pregnancies after PG		
7	3	2	37+	2.8	NA	No pregnancies after PG		
8	3	2	35+	3.3	NA	No pregnancies after PG		
9	4	2	27+	2.4	110	No pregnancies after PG		
10	3	2	22+	17	50	No pregnancies after PG		
11	3	2	25+	27	83	No pregnancies after PG		

<sup>a</sup>Collagen XVII (BP180)-enzyme-linked immunoassay (ELISA), U/ml, normal reference <9 U/ml.

GA: gestational age, weeks; NA: not available.