

Table SI. Demographics of cutaneous T-cell lymphoma-mycosis fungoides patients for immunohistochemistry experiments

Age, years/ Sex	Race	Anatomic region	Grouped stage <sup>a</sup>	Known pruritus	Slide staining	Free floating sections
50/M	C	Trunk	Early	Yes	Yes	Yes
48/M	C	Extremity	Early	Yes <sup>b</sup>	Yes	Yes
54/M	C	Trunk	Early	Unknown	Yes	Yes
77/F	C	Extremity	Later	Yes <sup>b</sup>	Yes	Yes
68/M	N/A	Trunk	Early	N/A	Yes	Yes
78/M	C	Trunk	Early	N/A	Yes	Yes
39/F	AA	Extremity	Early	Unknown	Yes	Yes
63/M	N/A	Trunk	Early	N/A	Yes	Yes
60/M	C	Trunk	Later	Yes	Yes	Yes
56/F	N/A	Trunk	Early	N/A	Yes	No
72/M	C	Trunk	Later	Yes	Yes	No
72/F	C	Head/neck/scalp	Early	Yes <sup>b</sup>	Yes	No
68/M	AA	Trunk	Later	Yes <sup>b</sup>	Yes	No
88/M	C	Trunk	Early	N/A	Yes	No
63/M	C	Trunk	Early	Yes <sup>b</sup>	Yes	No
73/M	C	Extremity	Early	Unknown	Yes	No
35/M	CV	Extremity	Early	Unknown	Yes	No

<sup>a</sup>Early: Stages IA and IB; Later: Stages IIA-IVB. <sup>b</sup>Pruritus documented to be present from lesion biopsied.

C: Caucasian; N/A: No chart available; AA: African-American; CV, Cape Verdian.