

Table SII. Clinical outcomes, treatment, follow-up and course of patients with ichthyosiform mycosis fungoides

Patient	Treatment and response	Follow-up duration after diagnosis, years	Course
1	PUVA → PR, 6 years later relapse, NBUVB → CR	18	No new lesions appeared
2	PUVA → PR	8.2	Responded to treatment but the lesions remained
3	PUVA, Aci → PR	10.8	Responded well after therapy, but lesions remained
4	PUVA, Aci → PR, MTX (for 31 months), Ste → PR	3	Responded to treatment but the lesions remained
5	PUVA, Aci → PR, MTX → PR, UVA1, Ste → PR	8	Responded to treatment but the lesions remained
6	PUVA, Aci → PR, Aci → PR	5	Responded to treatment but the lesions remained
7	PUVA → PR	1	Repigmentation occurred after PUVA, but the lesions remained
8	NBUVB → no response, PUVA, Aci → PR, PCALCL → Excision, IFN- $\alpha$ → PR	5	Not responded well after NBUVB, but combined treatment with PUVA and Aci showed partial remission
9	PUVA → PR	1	Follicular lesions and newly emerging nodules also appeared
10	NBUVB → PR	0.5	Responded well after therapy, but lesions remained

PUVA: psoralen plus ultraviolet A; NBUVB: narrowband ultraviolet B; UVA1: ultraviolet A1; MTX: methotrexate; Aci: oral acitretin; ste: topical steroid; CR: complete remission; PCALCL: primary cutaneous anaplastic large cell lymphoma; PR: partial remission; IFN- $\alpha$ : interferon  $\alpha$ .