

Table SI. Main clinical and histological characteristics of patients with and without telaprevir-skin eruptions

Case	Age, years/ Sex	Onset of eruption (week)	Clinical pattern	Localization of skin lesions	Severity	Histological examination	Suspension of antiviral treatment due to skin eruption
1	53/M	11	Eczematous	Head	Mild	Superficial perivascular predominantly lymphocytic infiltrate	–
2	35/F	Unknown	Eczematous	Unknown	Mild	–	–
3	60/M	9	Eczematous	Head, trunk, upper limbs and lower limbs	Moderate	Epidermal spongiosis and superficial perivascular predominantly lymphocytic infiltrate	–
4	59/M	8	Urticarial	Head and trunk	Moderate	–	–
5	58/F	8	Maculopapular	Trunk, upper limbs and lower limbs	Severe	Epidermal spongiosis and superficial perivascular predominantly lymphocytic infiltrate	TPV
6	46/M	12	Eczematous	Trunk, upper limbs and lower limbs	Mild	Superficial perivascular predominantly lymphocytic infiltrate	–
7	53/M	2	Urticarial	Trunk, upper limbs and lower limbs	Moderate	Discrete superficial perivascular predominantly lymphocytic infiltrate	–
8	52/F	8	Urticarial	Head, trunk, upper limbs and lower limbs	Severe	Superficial perivascular predominantly lymphocytic infiltrate and mild oedema in papillary dermis	TPV
9	52/M	5	Eczematous	Upper limbs	Mild	–	–
10	20/F	24	Eczematous	Trunk	Mild	–	–
11	56/M	28	Eczematous	Trunk	Mild	Discrete epidermal spongiosis and superficial perivascular predominantly lymphocytic infiltrate	–
12	49/M	Unknown	Eczematous	Trunk and lower limbs	Mild	–	–
13 ^a	53/M	8	DRESS	Head, trunk, upper limbs and lower limbs	Severe	Discrete epidermal spongiosis and superficial perivascular predominantly lymphocytic infiltrate	PEG-INF, RBV and TPV
14	41/M	20	Eczematous	Trunk, upper limbs and lower limbs	Moderate	–	–
15	62/F	1	Lichenoid dermatitis	Trunk, upper limbs and lower limbs	Severe	Superficial perivascular predominantly lymphocytic infiltrate with abundant eosinophils	TPV
16	52/M	Unknown	Eczematous	Unknown	Mild	–	–
17	64/F	12	Eczematous	Head, upper limbs and lower limbs	Mild	–	–
18	37/F	9	Eczematous (with vesicles)	Head, trunk, upper limbs and lower limbs	Severe	Epidermal spongiosis (with vesiculation) and superficial perivascular predominantly lymphocytic infiltrate and mild oedema in papillary dermis	PEG-INF, RBV and TPV
19	51/F	2	Maculopapular	Trunk and lower limbs	Moderate	–	–
20	53/F	11	Maculopapular	Trunk, upper limbs and lower limbs	Moderate	–	–
21	57/M	5	Maculopapular	Trunk, upper limbs and lower limbs	Severe	Not available	–
22	54/M	3	Erythema multiforme-like	Head, trunk, upper limbs and lower limbs	Severe	Superficial perivascular predominantly lymphocytic infiltrate	–
23	56/M	5	Eczematous	Head, trunk, upper limbs and lower limbs	Mild	–	–
24	49/M	10	Maculopapular	Trunk, upper limbs and lower limbs	Severe	Epidermal spongiosis and superficial perivascular predominantly lymphocytic infiltrate with abundant eosinophils	TPV
25	46/M	7	Eczematous	Lower limbs	Mild	–	–
26	62/F	Unknown	Eczematous	Unknown	Moderate	–	–
27	39/M	10	Eczematous	Head, trunk, upper limbs and lower limbs	Moderate	Epidermal spongiosis and superficial perivascular predominantly lymphocytic infiltrate with exocytosis	–
28	66/F	Unknown	Eczematous	Unknown	Mild	–	–
29	45/F	Unknown	Eczematous	Unknown	Mild	–	–

^aSee Martín Domínguez et al. (14).

DRESS: drug reaction with eosinophilia and systemic symptoms; TPV: telaprevir; PEG-INF: pegylated interferon; RBV: ribavirin.