

Table SIII. Summary of demographic and clinical characteristics for the patients who died after discharge

Pat No./Sex/ Age, years	Follow-up period <sup>a</sup>	Underlying disease	Sequelae	Cause of death	Death related to the sequelae of SJS/TEN
1/M/73	8.5 years	Cervical spondylosis	Chronic eczema	Unknown	No
2/M/77	3.5 years	HTN, COPD, Colon cancer	None	Lung cancer with multiple metastases	No
3/F/46	1.8 years	Breast cancer with multiple metastases	Dry eye syndrome Mebomian gland obstruction	Progression of breast cancer Urinary tract infection, sepsis	No
4/M/82	1.2 years	Pulmonary TB, CRI, hyperuricemia	Chronic pruritus	Pneumonia with respiratory failure	No
5/F/64	2.8 years	DM, HTN, Psoriasis	Sjögren's syndrome	Pneumonia with respiratory failure	No
6/F/77	3.3 years	DM, HTN, RA, PAOD	None	Acute upper airway obstruction	No
7/M/65	10.5 years	Psoriasis	None	Organophosphate intoxication	No
8/M/61	4.8 years	MEN type I CRI, gout	Chronic eczema	Acute subdural haemorrhage	No
9/F/78	1.2 years	DM, HTN	Generalized asteatotic eczema	Unknown primary neuroendocrine tumour with liver and lymph node metastasis	No
10/F/53	1.2 years	Breast cancer with brain metastasis	Dry eye syndrome	Unknown	No
11/M/71	1.0 years	Seizure, ESRD, CAD, AS, DM, HTN	None	Pneumonia with sepsis	No
12/M/76	4.0 months	Common bile duct and gall stone	Chronic conjunctivitis	Acute cholangitis with sepsis	No
13/39/M	9.9 months	Mitochondrial encephalopathy DM	None	Aspiration pneumonia with sepsis	No
14/87/F	2.6 months	HTN, old stroke	Generalized asteatotic eczema	Aspiration pneumonia with sepsis	No
15/80/M	6.8 months	Pulmonary TB Lymphoma	None	<i>Pneumocystis jiroveci</i> pneumonia	No

<sup>a</sup>Time from presentation with SJS/TEN to death.

AS: ankylosing spondylitis; CAD: coronary artery disease; COPD: chronic obstructive pulmonary disease; CRI: chronic renal insufficiency; DM: diabetes mellitus; ESRD: end-stage renal disease; F: female; HTN: hypertension; M: male; MEN: multiple endocrine neoplasia; RA: rheumatoid arthritis; TB: tuberculosis.