

Table SI. Characteristics of mycosis fungoides patients receiving systemic treatment and chemotherapy, and the choice of drugs

Age at diagnosis, years	Sex	Initial stage	TNMB	Highest stage	Immunomodulating therapy	Chemotherapy	Other systemic treatment	Deceased within study period
46	M	IA	T1a	IA	Acitretin			No
49	M	IB	T2b	IIA	Acitretin, Isotretinoin, Interferon, Bexarotene, Prednisone	Mtx, Chlorambucil, pegylated liposomal doxorubicin	Alemtuzumab	No
50	M	IIB	T3N1a <sup>a</sup>	IIB <sup>b</sup>		CHOP		No
53	M	IA	T1b <sup>a</sup>	IIIA <sup>b</sup>	Bexarotene	Mtx, CHOP, Gemcitabine, pegylated liposomal doxorubicin		No
54	M	IB	T2b <sup>a</sup>	IVB	Isotretinoin, Interferon	Mtx, CHOP, Gemcitabine	Alemtuzumab	Yes <sup>c</sup>
54	F	IA	T1a	IB	Acitretin			No
58	M	IA	T1a	IA		Mtx		No
63	M	IA	T1b <sup>a</sup>	IA	Isotretinoin			No
64	F	IIB	T3N1a	IIB		Mtx		No
65	F	IIA	T1bN2b	IIA <sup>b</sup>	Bexarotene	CHOP, Gemcitabine		No
65	M	IB	T2b <sup>a</sup>	IIB	Prednisolone	Mtx		Yes <sup>c</sup>
69	F	IB	T2b	IVA2	Isotretinoin, Prednisone	Mtx, Chlorambucil	Alemtuzumab	Yes <sup>c</sup>
71	M	IA	T1b <sup>a</sup>	IVB <sup>b</sup>	Acitretin, Isotretinoin, Prednisolone	Gemcitabine, Chlorambucil, CHOP	Alemtuzumab	Yes
75	M	IIA	T2bN1a	IIIA	Prednisone	Chlorambucil		Yes <sup>c</sup>
75	F	IA	T1b <sup>a</sup>	IA	Isotretinoin			Yes
76	F	IB	T2b	IIB	Acitretin	Mtx		Yes
83	M	IIIA	T4N0	IIIA	Prednisolone	Mtx		No
84	M	IB	T2N0B1b <sup>a</sup>	IB		Mtx		No

<sup>a</sup>Adnexal disease. <sup>b</sup>Large cell transformation. <sup>c</sup>Disease-specific mortality.  
Mtx: methotrexate; CHOP: cyclophosphamide, doxorubicin, vincristine, prednisone.