

Table SI. Values for variables (total naevi count, the ratio between trunk naevus count over total naevus count, and the ratio between right hemisoma naevus count over total naevus count) for the variables considered (n = 111)

	n (%)	p-value	Total naevi count Mean ± SD	p-value	Trunk and head naevus count/total naevus count Mean ± SD	p-value	Right hemisoma naevus count/total naevus count Mean ± SD	p-value
Sex								
Male	52 (46.8)		30 ± 5		0.62 ± 0.1		0.54 ± 0.1	
Female	59 (53.2)	NS	31 ± 5	NS	0.55 ± 0.2	<0.01	0.52 ± 0.1	NS
Age, (mean ± SD: 44.1 ± 16.3)								
<44 years	64 (58)		34 ± 5		0.60 ± 0.2		0.52 ± 0.1	
>45 years	47 (42)	NS	26 ± 5	<0.01	0.56 ± 0.2	NS	0.53 ± 0.1	NS
BMI, kg/m ² (mean ± SD 24 ± 4.4)								
≤24	61 (55)		31 ± 5		0.56 ± 0.1		0.53 ± 0.1	
≥25	50 (45)	NS	29 ± 5	NS	0.60 ± 0.2	NS	0.53 ± 0.1	NS
Cigarette smoking								
Smokers	32 (28.8)		33 ± 8		0.58 ± 0.2		0.51 ± 0.1	
Not smokers	79 (71.2)	<0.01	30 ± 4	NS	0.58 ± 0.2	NS	0.53 ± 0.1	NS
Having parents with numerous moles								
Positive	38 (34.2)		49 ± 8		0.60 ± 0.2		0.52 ± 0.1	
Negative	73 (65.8)	<0.01	26 ± 4	<0.01	0.57 ± 0.2	NS	0.53 ± 0.1	NS
Childhood sun exposure (mean ± SD 4.6 ± 2.4, range: 1–9)								
Low	55 (49.5)		31 ± 5		0.59 ± 0.2		0.53 ± 0.1	
High	56 (50.5)	NS	30 ± 6	NS	0.57 ± 0.1	NS	0.53 ± 0.1	NS
Adulthood sun exposure (mean ± SD 19.4 ± 5.3, range: 1–33)								
Low	47 (42)		34 ± 5		0.59 ± 0.2		0.53 ± 0.1	
High	64 (58)	NS	30 ± 5	NS	0.57 ± 0.1	NS	0.53 ± 0.1	NS
Skin phototype								
Fair (≤II)	40 (36)		25 ± 7		0.54 ± 0.2		0.5 ± 0.1	
Dark (>II)	71 (64)	<0.01	32 ± 4	<0.05	0.60 ± 0.3	<0.05	0.51 ± 0.1	<0.05
Skin photodamage								
Low	69 (62.2)		26 ± 4		0.59 ± 0.2		0.52 ± 0.1	
High	42 (37.8)	<0.05	43 ± 8	<0.01	0.56 ± 0.2	NS	0.53 ± 0.1	NS

SD: standard deviation; BMI: body mass index; NS: statistically not significant.