

Table SIII. Demodex densities in patients with seborrhoeic dermatitis or acne vulgaris according to the associated dermatoses: the Dd was high only if there was associated demodicosis ($p < 0.001$)

	Patients n (%)	SSSB 1 Mean \pm SEM	SSSB 2 Mean \pm SEM	SSSB 1+2 Mean \pm SEM
<i>Seborrhoeic dermatitis</i>				
No other dermatoses	33 (19.0)	0.5 \pm 0.2	1.8 \pm 0.5	2.2 \pm 0.6
Demodicosis only	134 (77.0)	74.6 \pm 8.2	152.3 \pm 13.8	226.9 \pm 19.5
Other dermatosis (not demodicosis)	7 (4.0)	0.1 \pm 1.4	0.7 \pm 0.7	0.9 \pm 0.9
Seborrhoeic dermatitis, All ^a	174 (100.0)	60.4 \pm 6.8	119.0 \pm 11.1	179.4 \pm 16.2
<i>Acne vulgaris</i>				
No other dermatoses	70 (64.2)	0.3 \pm 0.1	0.9 \pm 0.3	1.2 \pm 0.4
Demodicosis only	31 (28.4)	89.0 \pm 23.5	175.2 \pm 33.1	264.2 \pm 48.3
Other dermatosis (not demodicosis)	8 (7.3)	0.4 \pm 0.2	2.6 \pm 1.5	3.0 \pm 1.7
Acne vulgaris, all ^b	109 (100.0)	25.52 \pm 7.6	50.6 \pm 12.0	76.14 \pm 17.7

^aFour patients with seborrhoeic dermatitis and demodicosis and another dermatosis were not considered in this Table. ^bTwo patients with acne vulgaris and demodicosis and another dermatosis were not considered in this Table. SEM: standard error of the mean; SSSB: standardized skin surface biopsy.