

Table SII. Demographic findings of participants in this study according to psychiatric visits

	Psychiatric visit			p-value
	No n (%)	Yes n (%)	Total n	
Sex				
Male	179,965 (93.3)	12,962 (6.7)	192,927	< 0.001
Female	166,111 (93.8)	10,981 (6.2)	177,092	
Enrolled age*				
0 years	104,492 (96.3)	4,018 (3.7)	108,510	< 0.001
1–5 years	46,995 (92.2)	3,967 (7.8)	50,962	
6–10 years	67,307 (92.7)	5,335 (7.3)	72,642	
10–14 years	64,224 (92.4)	5,291 (7.6)	69,515	
15–20 years	63,058 (92.2)	5,332 (7.8)	68,390	
Residential area				
Rural	107,423 (94.3)	6,484 (5.7)	113,907	< 0.001
Urban	238,653 (93.2)	17,459 (6.8)	256,112	
Type of insurance**				
Employee	150,693 (93.2)	11,060 (6.8)	161,753	< 0.001
Self-employed	186,573 (93.8)	12,232 (6.2)	198,805	
Medical aid	8,810 (93.1)	651 (6.9)	9,461	
Income level				
Low	35,220 (93.2)	2,588 (6.8)	37,808	< 0.001
Low–middle	78,891 (93.6)	5,384 (6.4)	84,275	
Middle–high	141,274 (94.0)	9,013 (6.0)	150,287	
High	90,691 (92.9)	6,958 (7.1)	97,649	
Alopecia areata				
No	341,852 (93.6)	23,460 (6.4)	365,312	
Alopecia areata	3,917 (89.8)	444 (10.2)	4,361	
Alopecia totalis	173 (88.7)	22 (11.3)	195	
Alopecia universalis	134 (88.7)	17 (11.3)	151	
Total	346,076 (93.5)	23,943 (6.5)	370,019	