

**Table SI. Baseline demographics of patients enrolled in the pooled FIXTURE and ERASURE studies who completed the Psoriasis Symptom Diary at baseline (n = 820)**

Characteristics	Secukinumab 150 mg (n = 229)	Secukinumab 300 mg (n = 226)	Etanercept 50 mg (n = 137)	Placebo (n = 228)
Age, mean ± SD, years	45.7 ± 12.9	43.3 ± 13.4	43.7 ± 13.1	43.1 ± 11.7
Male, n (%)	151 (65.9)	142 (62.8)	98 (71.5)	163 (71.5)
White, n (%)	190 (83.0)	189 (83.6)	107 (78.1)	185 (81.1)
Body weight, kg, mean ± SD	89.3 ± 22.2	90.2 ± 23.1	87.8 ± 21.7	89.4 ± 23.2
Body mass index, kg/m <sup>2</sup> , mean ± SD	29.8 ± 6.8	30.4 ± 7.4 <sup>a</sup>	29.4 ± 6.5	29.7 ± 7.3
PASI (0–72), mean ± SD (range)	21.8 ± 9.0 (12.0–69.6)	21.9 ± 9.0 (12.0–64.2)	21.3 ± 8.0 (12.0–49.6)	21.7 ± 8.9 <sup>b</sup> (10.6–55.5)
Body surface area, %, mean ± SD	31.2 ± 18.0	29.2 ± 17.6	29.7 ± 16.9	29.0 ± 15.6
Investigator's Global Assessment (0–4) category, n (%)				
Moderate (IGA = 3)	152 (66.4)	143 (63.3)	87 (63.5)	139 (61.0)
Severe (IGA = 4)	77 (33.6)	83 (36.7)	50 (36.5)	89 (39.0)
Psoriasis Symptom Diary scores (0–10) <sup>c</sup> , mean ± SD				
Itching severity	6.46 ± 2.37	6.42 ± 2.39	6.18 ± 2.43	6.06 ± 2.45
Pain severity	5.29 ± 3.06	5.49 ± 2.99	5.28 ± 2.96	4.97 ± 3.03
Scaling severity	6.47 ± 2.41	6.40 ± 2.55	6.33 ± 2.61	6.18 ± 2.44

<sup>a</sup>n = 225. <sup>b</sup>n = 227. <sup>c</sup>A score of 0 = no itching/pain/scaling; 10 = itching/pain/scaling as bad as you can imagine. SD: standard deviation; PASI: Psoriasis Area and Severity Index.