

Table SII. Multiple linear regression analysis evaluating the association between basal and squamous cell carcinoma quality of life (BaSQoL) subscales and satisfaction with information (* $p < 0.05$)

	Behaviour		Diagnosis and treatment		Worries		Appearance		Other people	
	Beta	95% CI	Beta	95% CI	Beta	95% CI	Beta	95% CI	Beta	95% CI
Satisfaction with information (dissatisfied = ref)										
Satisfied	-0.15*	-0.25/-0.05	-0.12*	-0.25/-0.00	-0.25*	-0.35/-0.15	-0.14*	-0.20/-0.08	-0.07	-0.18/0.05
Age at time of questionnaire	-0.01*	-0.01/-0.00	-0.01*	-0.02/-0.00	-0.01*	-0.02/-0.00	-0.01*	-0.01/-0.01	-0.01*	-0.01/-0.00
Time since diagnosis	-0.01	-0.04/0.03	0.07*	0.02/0.12	0.07*	0.02/0.12	0.06*	0.02/0.09	-0.02	-0.5/0.08
Sex (male = ref)										
Female	0.07	-0.02/0.16	0.29*	0.17/0.40	0.12*	0.02/0.21	0.02	-0.04/0.07	0.20*	0.09/0.30
Education (low = ref)										
Medium	-0.14*	-0.24/-0.04	-0.03	-0.16/0.10	-0.04	-0.14/0.06	-0.04	-0.10/0.02	0.13*	0.01/0.25
High	0.20*	0.02/0.38	-0.15	-0.38/0.08	-0.15	-0.33/0.03	-0.08	-0.19/0.02	-0.06	-0.16/0.28
Occupation ((self-) employed = ref)										
Unemployed	-0.08	-0.22/0.06	-0.14	-0.32/0.03	-0.05	-0.19/0.09	0.08	-0.00/0.16	-0.02	-0.18/0.15
Partnership										
No partner (partner = ref)	0.08	-0.03/ 0.19	0.12	-0.02/0.26	-0.07	-0.18/0.04	0.05	-0.01/0.12	-0.00	-0.13/0.13
Location of tumour (face = ref)										
Other	-0.11*	-0.21/-0.01	-0.14*	-0.26/-0.01	0.02	-0.09/0.12	-0.09*	-0.15/-0.03	0.07	-0.05/0.20
Type of tumour (BCC= ref)										
SCC	0.04	-0.11/0.18	0.14	-0.05/0.32	0.29*	0.14/0.44	0.05	-0.04/0.13	0.25*	0.07/0.43
Comorbidity (no = ref)										
1	-0.08	-0.22/0.06	0.09	-0.09/0.27	0.01	-0.13/0.15	0.07	-0.01/0.15	0.03	-0.14/0.20
≥2	-0.02	-0.13/0.08	0.11	-0.02/0.25	0.04	-0.07/0.15	0.07*	0.00/0.13	0.13	0.00/0.25

CI: confidence interval; BCC: basal cell carcinoma; SCC: squamous cell carcinoma.