

Table SII. Treatment of the intertrigo

Orientation (<i>n</i> =62)	
Hospitalization, <i>n</i> (%)	47 (75.8)
Length of stay, median in days (range)	11 (3-33)
Initial, <i>n</i> (%)	18 (29)
After outpatient treatment fail, <i>n</i> (%)	29 (46.8)
Outpatients treatment, <i>n</i> (%)	15 (24.2)
Duration of outpatient follow-up, weeks, median (range) (3 NAD)	7 (1-40)
Care performed by a nurse, <i>n</i> (%)	5
Number of consultations, median (range)	2 (0-10)
Local treatment alone (<i>n</i> =62), <i>n</i> (%)	34 (54.8)
Nurse at domicile, <i>n</i> (%)	40 (64.5)
Material used to separate toes, <i>n</i> (%)	50 (80.6)
Detersion, <i>n</i> (%)	19 (30.6)
Salicylic acid	9 (14.5)
Mecanic debridment	10 (16.1)
Antiseptics, <i>n</i> (%)	51 (82.3)
One	35
Two	14
Three	2
Hydrogen peroxide	17
Boric acid	15
Chlorhexidine	8
Sodium Hypochorite	8
Povidone iodine	7
Silver nitrate	6
Potassium permanganate	3
Acetic acid	3
Eosine	2
Local antibiotics, <i>n</i> (%)	27 (43.5)
Silver sulfadiazine	16
Gentamycin	7
Other (colymicine, metronidazole, fusidic acid)	4
Systemic antibiotics (<i>n</i> =28), <i>n</i>	
For intertrigo	25
For chondritis (PA)	1
For urinary infection	2
Modalities	
Per Os	13
Intravenous	15
Antibiotics used	
Against PA (<i>n</i> =16)	10
Monotherapy	
Piperacillin-tazobactam	5
Ceftazidime	2
Ciprofloxacin	2
Oflloxacin	1
Bitherapy	
Piperacillin-tazobactam + ciprofloxacin	6
Imipenem + ciprofloxacin	4
Ceftazidime + ciprofloxacin	1
Against SA/SBHA (12)	1
Pristinamycin	5
Amoxicillin-clavulanic acid	4
Amoxicillin	1
Linezolid	1
Clindamycin	1
Duration, days, median (range)	10 (3-21)

NAD: non available data.