

Table SI. Mean minimal clinically important difference (MCID) for visit 2 vs. visit 1 (V1-V2) and for visit 3 vs. visit 1 (V1-V3) per average pruritus severity grade in the 24 h preceding the corresponding visit as rated using a visual analogue scale (VAS_{24h}) and numeric rating scale (NRS_{24h})

Pruritus severity grade	VAS _{24h}						NRS _{24h}					
	<i>n</i>	Min	Max	Median	Mean	SD	<i>n</i>	Min	Max	Median	Mean	SD
MCID V1-V2												
1: 0-2.9	13	-4.50	1.00	0.00	-0.62	1.72	9	-4.00	2.00	0.00	0.00	1.87
2: 3-6.9	33	-2.50	4.50	1.00	0.83	1.83	34	-3.00	3.00	1.00	0.46	1.54
3: 7-8.9	13	0.50	6.00	2.50	2.85	1.52	19	0.00	6.00	3.00	2.74	1.52
4: 9-10	9	1.50	7.00	4.00	4.33	2.15	7	2.00	7.00	4.00	4.21	1.91
MCID V1-V3												
1: 0-2.9	12	-2.50	1.00	0.25	-0.04	1.12	9	-2.00	2.00	0.00	0.22	1.39
2: 3-6.9	32	-3.00	5.00	1.50	1.34	1.94	33	-4.00	3.00	2.00	1.06	1.82
3: 7-8.9	13	-1.00	6.00	4.00	3.65	2.00	18	-1.00	7.00	3.00	2.92	2.12
4: 9-10	9	-0.50	8.50	4.50	4.56	2.93	8	0.00	8.00	4.50	4.38	2.50

SD: standard deviation.