

Table SII. Mean production values of cytokines in lesional skin of adult and paediatric psoriasis patients, according to psoriasis type

Cytokines	Plaque psoriasis			Guttate psoriasis			Pustular psoriasis		
	Adult patients (n = 17)	Paediatric patients (n = 15)	p-value	Adult patients (n = 6)	Paediatric patients (n = 6)	p-value	Adult patients (n = 2)	Paediatric patients (n = 1)	p-value
Qualitative analysis, (median, Grade 0-3)									
CD3	1+	1+	0.551	1+	1+	0.589	1.5+	1+	0.667
CD4	1.5+	2+	0.502	1.25+	1+	0.485	2+	0+	0.667
CD8	2+	3+	0.002	1.25+	3+	0.132	1.5+	3+	1.000
CD45	1+	1+	0.153	1.5+	1+	0.310	1.5+	0+	0.667
TNF- α	2+	3+	<0.001	0+	2+	0.002	1.75+	3+	1.000
FoxP3	0.5+	0+	0.002	0.5+	0+	0.065	1.5+	0+	0.667
IL-13a	1+	0.5+	0.390	0.75+	0.5+	0.589	1.25+	1+	1.000
IL-17a	2+	0.5+	0.000	2+	0.25+	0.065	2+	1+	0.667
IL-22	1+	0.5+	0.165	1+	1+	0.699	0.75+	0.5+	0.667
IFN- γ	2+	2+	0.766	2.75+	2.25+	0.699	2+	2+	1.000
Quantitative analysis, (median, % of stained cells in the whole layer)									
TNF- α	1.97	11.3	<0.001	1.14 ^a	7.18 ^a	0.012	4.06	1.91	1.000
IL-13 ^a	6.09 ^a	5.30 ^a	0.505	5.76 ^a	7.17 ^a	0.374	2.66	4.02	1.000
IL-17 ^a	6.91	2.60	0.058	8.24	3.18	0.180	13.6	31.1	1.000
IL-22	13.3	11.5	0.710	11.4 ^a	16.2 ^a	0.616	10.0	4.51	0.667
IFN- γ	33.2 ^a	30.2 ^a	0.338	24.9 ^a	29.8 ^a	0.954	22.5	39.3	1.000

^aMean value. Significant values are shown in bold.

TNF: tumour necrosis factor; IL: interleukin; IFN: interferon.