

Table SI. Main characteristics of previous cases of diffuse dermal angiomatosis with multifocal involvement

Author	Sex	Age, years	Medical history	Affected locations	Treatment	Evolution
Krell et al. (1)	F	47	PAD	Leg and thigh	Revasc	Improvement
	F	63	PAD, axillary bifemoral bypass graft, cardiac arrhythmia, smoking	Left and upper medial thigh and groin	Revasc	Significant improvement
Tollefson et al. (4)	F	51	Takeyashu arteritis with occlusion of both subclavian arteries, HBP, smoking	Breasts, thighs	Revasc	Significant improvement
Steele et al. (5)	M	72	CRF secondary to sarcoidosis on daily peritoneal dial, calciphylaxis	Thighs, calf, lower abdomen	Sodium thiosulphate	Death 4 weeks after the diagnosis
Halbesleben et al. (6)	M	38	HBP, cutis marmorata telangiectatica congenita	Arm, forearm, hand, posterior shoulder	Surgical removal of a painful plaque on the arm	No recurrence during 6-month follow-up
Yang et al. (7)	F	49	Polysubstance abuse, smoking, hepatitis C	Right breast, flank and hip	NS	NS
Gianotti et al. (8)	F	72	HBP, atrial fibrillation, thyroid cancer	Lower abdomen, thighs and buttocks	Methylprednisolone Warfarin withdrawal	No improvement, death after a few days
Crickx et al. (9)	F	71	PAD with lower-limb amputation, coronary by-pass surgery, diabetes mellitus, HBP, CRF, atrial fibrillation, ex-smoker	Trunk, limbs	Colchicine, prednisone, oxygen, stopping warfarin, controlled blood pressure and diabetes	Death 1 month later
Garcia-Colmenero et al. (10)	F	63	HBP, dyslipidaemia, smoking	Thighs, lower abdomen, perianal area	Control of cardiovascular risk factors, acetylsalicylic acid 100 mg/day, Revasc	Significant improvement
Ormerod et al. (11)	F	54	CRF, diabetes mellitus, PAD, cerebrovascular disease, parathyroidectomy	Forearm, abdominal wall	Removal of the arterio-venous fistula	Significant improvement
Mayor-Ibarguren et al. (12)	F	60	Liposarcoma treated with trabectedin and pegfilgrastim	Extensor surfaces of the lower extremities, elbows, back of the hands, pavilion of ears	Trabectedin and pegfilgrastim withdrawal	Death 6 months later after surgical complications and renal failure
Kirkland et al. (13)	F	58	HBP, PAD, bilateral iliac stents, smoking	Thigh, buttock, lower abdomen	Failure of Revasc.	Death of urosepsis 1 month later
Kimyai-Asadi et al. (14)	F	57	HBP, PAD, CRF, smoking	Both thighs, ankle	Revasc	Significant improvement

NS: not specified; PAD: peripheral arterial disease; CRF: chronic renal failure; Revasc: revascularization, HBP: high blood pressure.