

Table SVI. Summary of Kaplan–Meier overall survival estimates and follow-up

Overall survival analysis Ref, <i>n</i>	Total 207	Non-Metastatic 125	Metastatic 82	Metastatic 82	Metastatic 82
Censored overall, <i>n</i> (%)	46 (22.2)	24 (19.2)		22 (26.8)	
Dead overall, <i>n</i> (%)	161 (77.8)	101 (80.8)		60 (73.2)	
Follow-up time, months					
Median (IQR)	47.0 (20–98)	64.0 (25–119)		32.0 (17–67)	
Mean	66.5	74.0		55.1	
Range	0–277	1–254		0–277	
Overall survival	From initial diagnosis of first cSCC		From initial diagnosis of first primary mcSCC		From initial diagnosis of first metastasis
1-year survival percentage, estimate	92.8	92.8	92.7	92.7	68.0
Censored within first year, <i>n</i> (%)	1 (0.5)	0 (0.0)	1 (1.2)	1 (1.2)	2 (2.4)
Dead within first year, <i>n</i> (%)	15 (7.2)	9 (7.2)	6 (7.3)	6 (7.3)	26 (31.7)
2-year survival percentage, estimate	71.7	76.8	66.5	63.8	43.4
Censored within 2 years, <i>n</i> (%)	3 (1.9)	0 (0.0)	3 (3.7)	4 (4.7)	5 (6.1)
Dead within 2 years, <i>n</i> (%)	56 (27.1)	29 (23.2)	27 (32.9)	29 (35.4)	45 (54.9)
3-year survival percentage, estimate	62.1	66.4	54.8	50.5	31.6
Censored within 3 years, <i>n</i> (%)	6 (3.9)	0 (0.0)	6 (7.3)	7 (8.5)	9 (11.0)
Dead within 3 years, <i>n</i> (%)	78 (37.2)	42 (33.6)	36 (43.9)	39 (47.6)	53 (64.6)
4-year survival percentage, estimate	53.9	60.0	43.6	38.9	31.6
Censored within 4 years, <i>n</i> (%)	10 (6.3)	0 (0.0)	10 (12.2)	11 (13.4)	13 (15.9)
Dead within 4 years, <i>n</i> (%)	94 (44.9)	50 (40.0)	44 (53.7)	47 (57.3)	53 (64.6)
5-year survival percentage, estimate	46.4	52.0	37.2	30.7	31.6
Censored within 5 years, <i>n</i> (%)	12 (6.3)	0 (0.0)	12 (14.6)	13 (15.9)	14 (17.1)
Dead within 5 years, <i>n</i> (%)	108 (51.7)	60 (48.0)	48 (58.5)	52 (63.4)	53 (64.6)
<i>p</i> -value ^a			0.03	0.002	<0.001

^aCompared with non-metastatic patient cohort.

IQR: interquartile range; cSCC: cutaneous squamous cell carcinoma; mcSCC: metastatic cutaneous squamous cell carcinoma.