

Table SII. Melanoma-related characteristics in patients with mycosis fungoides (MF) and psoriasis

Characteristics	MF (n = 47)	Psoriasis (n = 23)	p-value
Site of melanoma, n (%)			0.388
Head and neck	10 (23.3)	2 (10)	
Limbs	12 (27.9)	8 (40)	
Trunk	21 (48.8)	10 (50)	
Two areas	1	0	
Mucosal	1	1	
Unknown	2	2	
Multiple melanoma (≥ 2), n	5	2	1.0
Type of melanoma, n			N/A
SSM	17	4	
LM	2	0	
LMM	2	0	
Nodular	0	1	
SSM+LM	1	0	
Unknown	25	18	
Depth, n (%)			0.005
<i>In situ</i>	28 (63.6)	6 (30.0)	
Thin ≤ 1 mm	13 (29.5)	8 (40.0)	
Thick > 1 mm	3 (6.8)	6 (30.0)	
Unknown	3	3	
Breslow depth, mm mean ± SD	0.26 ± 0.4	1.30 ± 1.6	0.001

SD: standard deviation; SSM: superficial spreading melanoma; LM: lentigo maligna; NM: nodular melanoma; LMM: lentigo maligna melanoma; seq: sequential; N/A: not applicable.