

Table SII. Odds ratios (ORs) and 95% confidence intervals (95% CIs)^a for the association between phenotypic characteristics and melanoma T category by body site in the Norwegian Women and Cancer (NOWAC) cohort

Phenotypic characteristics	Trunk, <i>n</i> = 395		Lower limb, <i>n</i> = 446		Upper limb, <i>n</i> = 189		Head/neck, <i>n</i> = 84	
	Cases, <i>n</i> ^b	OR (95% CI)	Cases, <i>n</i> ^b	OR (95% CI)	Cases, <i>n</i> ^b	OR (95% CI)	Cases, <i>n</i> ^b	OR (95% CI)
Pigmentary score								
Dark	24	1.00	28	1.00	8	1.00	4	1.00
Medium	200	0.46 (0.19–1.12)	199	0.64 (0.26–1.60)	79	0.53 (0.08–3.36)	44	5.23 (0.23–116.2)
Fair	92	0.27 (0.10–0.70)	114	0.81 (0.31–2.10)	41	1.57 (0.22–11.11)	18	8.01 (0.32–198.2)
Small symmetrical naevi on arms/legs								
<50	156	1.00	180	1.00	89	1.00	39	1.00
≥51	53	0.79 (0.33–1.90)	59	1.69 (0.84–3.40)	20	5.11 (1.30–20.11)	10	3.04 (0.36–25.80)
Trend ^c		1.18 (0.78–1.78)		1.11 (0.75–1.63)		1.81 (1.07–3.05)		1.28 (0.54–3.03)
Small symmetrical naevi/pigmentary score								
< 50/Darker	117	1.00	125	1.00	54	1.00	31	– ^d
≥ 51/Darker	33	2.82 (0.58–13.68)	36	1.82 (0.31–10.68)	11	16.14 (0.53–495.01)	7	– ^d
< 50/Fairer	41	0.65 (0.28–1.51)	53	1.38 (0.45–4.19)	19	1.58 (0.30–8.35)	12	– ^d
≥ 51/Fairer	14	0.46 (0.14–1.45)	24	2.45 (0.72–8.30)	7	9.23 (1.21–70.34)	4	– ^d
<i>p</i> for interaction		0.15		0.97		0.60		– ^d

^aOrdinal logistic regression, adjusted for age at diagnosis, birth-cohort, education, place of residence, and subtype. ^bNumbers may not add up to the total due to missing.

^cEstimation of linear trend conducted by treating the variable "Small symmetrical naevi on arms/legs" (with categories 0, 1–10, 11–50, ≥51) as continuous in the model.

^dEstimation was not calculated, due to small sample size.