

Table SI. Associations between tumour size and other parameters

	Median tumour size mm (range)	Mean tumour size mm (95% CI)	p-value
Localization			
Nose	9.0 (3–35)	9.4 (8.5–10.4)	< 0.0001
Ear	10.0 (2–40)	11.4 (9.6–13.2)	
Scalp	12.0 (5–50)	16.2 (12.9–19.6)	
Periorbital	9.5 (2–22)	9.0 (7.2–10.8)	
Cheek, forehead, perioral and temple	10.0 (2–50)	11.4 (10.7–12.0)	
Trunk (including neck)	12.0 (4–200)	18.2 (14.4–22.0)	
Upper extremities	15.0 (5–60)	17.9 (14.1–21.8)	
Lower extremities	12.0 (4–80)	17.8 (12.7–22.9)	
Localization, subgrouped			
Face and scalp	10.0 (2–50)	11.3 (10.7–11.8)	< 0.0001
Other locations	12.0 (4–200)	18.1 (15.4–20.7)	
Physician specialty			
General practitioner	10.0 (4–35)	9.8 (8.4–11.3)	< 0.0001
General surgeon	13.0 (4–40)	14.9 (12.7–17.0)	
Plastic surgeon	12.0 (3–200)	17.0 (14.9–19.1)	
Dermatologist	10.0 (3–60)	11.1 (10.1–12.2)	
Otorhinolaryngologist	10.0 (2–35)	10.3 (9.6–11.0)	
Other	9.0 (4–40)	12.9 (5.1–20.7)	
Physician experience			
Specialist	10.0 (2–200)	13.1 (12.0–14.2)	0.41
Resident	10.0 (4–40)	12.9 (11.5–14.3)	

Significant p-values in bold.