

SHORT REPORTS

Effect of Retinoic Acid on the Infiltration of Murine Melanoma Cells into the Type I Collagen Gel

TAKESHI KONO, MASAYOSHI FURUKAWA, TSUKASA TANII, SHOJI TANIGUCHI, NOBUYUKI MIZUNO, MASAMITSU ISHII and TOSHIO HAMADA

Department of Dermatology, Osaka City University Medical School, Osaka, Japan

Two lines of murine melanoma cells (B16 and Cloudman S91) were cultured on type I collagen gel and the effects of all-trans-retinoic acid on the growth and infiltration into the gel were assayed. In both lines, proliferation and the degree of infiltration were suppressed by the addition of all-trans-retinoic acid. The infiltration-inhibiting effect was expressed very rapidly and was dose-dependent at concentrations ranging from 10^{-7} to 10^{-5} M of all-trans-retinoic acid. These results suggest the anti-invasive effects of all-trans-retinoic acid on melanoma cells.

(Accepted May 16, 1990.)

Acta Derm Venereol (Stockh) 1991; 71: 41-44.

T. Kono, Department of Dermatology, Osaka City University Medical School, 1-5-7, Asahimachi, Abeno-ku, Osaka 545, Japan.

Tumour cells tend to invade surrounding normal tissue by penetration or infiltration with active movement. However, the precise mechanism is not well elucidated because of the absence of simple *in vitro* experimental systems (1, 2).

We and other authors have reported that cells cultured in or on type I collagen gel resemble, as closely as possible, those *in vivo* (3-7). Recently, Schor et al. (8, 9) and our group (10-13) have cultured melanoma cells on and in type I collagen gel and have suggested melanoma cell-to-collagen interactions. Schor et al. first reported the penetration or infiltration phenomenon of tumour cells into collagen gel (8). On the other hand, all-trans-retinoic acid (RA) has been known to be an anti-proliferative agent against melanoma cells (14, 15).

In the present study, we cultured two lines of murine melanoma cells (B16 and Cloudman S91) on type I collagen gel and investigated the effects of RA on growth and infiltration into the gel.

MATERIALS AND METHODS

Cell culture

A B16 CBC7 clone was derived from B16 C₂M cells (16) (kindly donated by Dr A. Oikawa, Tohoku University School of Medicine, Sendai, Japan) and maintained in our laboratory. Cloudman S91 melanoma cells were obtained from the Dainippon Pharmaceutical Co. (Osaka, Japan). These lines were grown in Eagle's medium supplemented with 10% fetal calf serum (FCS).

Preparation of the type I collagen gel

Hydrated type I collagen gel was prepared using a modified method of Elsdale & Bard (3) as previously described (5-7, 10-13). Briefly, appropriate volumes of 0.3% pepsinized type I collagen solution (Nitta Gelatine Co., Osaka, Japan), concentrated Eagle's medium, 200 mM HEPES/0.08 N NaOH solution and FCS were mixed rapidly at a temperature below 4°C to a final collagen concentration of 2.0 mg/ml. One millilitre of the mixture was poured into a 35-mm bacteriological dish (Falcon) and incubated at 37°C. The collagen gelled within 10 min. Before use, cells were trypsinized, washed and then inoculated onto the gel at a cell density of 10^5 cells/dish.

Assay of total cell growth and infiltration into the collagen gel

The total number of cells both on and in the gel and that on the surface of the gel were determined by the modified method of Schor (8). The collagen gel with cells on its surface was washed three times with Ca²⁺, Mg²⁺-free phosphate-buffered saline (PBS²⁻) and incubated in 2 ml of 0.05% trypsin/2 mM EGTA in PBS²⁻ for 30 min. The trypsin solution was vigorously stirred by pipetting to release the cells. The cells released were recovered as completely as possible and stored in a tube. The gel was incubated in 2 ml of 0.4 mg/ml collagenase (Sigma) in serum-free medium for 20 min at 37°C. The collagenase solution was stirred vigorously by pipetting. The detached cells were recovered and added to the tube containing cells previously released by trypsinization and the cell numbers were counted using a hemocytometer (the number of cells on the surface of the gel; A). By the treatment with trypsin and collagenase, the cells on the surface were almost completely detached and the intact gel was left.

The gel was minced with scissors and incubated again in 2 ml of 0.4 mg/ml of collagenase for 5 h to dissolve the gel. Cells which infiltrated and grew in the gel were released,




Fig. 1. Time course of growth (A, B) and infiltration (C, D) of B16 (A, C) and S91 (B, D) melanoma cells. RA was added at a concentration of 10^{-6} M (○). Control, (●). Each point represents the mean of three experiments (\pm SD).

recovered and the cell numbers were counted (B). The total number of cells was the sum of A and B.

RA (Sigma) was added at the indicated concentration from day 0 of culturing. In the control, the same volume of solvent (ethanol) as that of RA was added.

Statistical analysis was performed using Student's *t*-test.

RESULTS

Growth-inhibitory effect of RA

Fig. 1 shows the time course of the growth (A, B) and the infiltration into the gel (C, D) of B16 (A, C) and S91 (B, D) melanoma cells. With the addition of RA (10^{-6} M), growth was suppressed in both lines. The suppression was shown to be stronger in S91 than in B16 cells (% of growth inhibition at day 5 was about 58% in B16 and 81% in S91 cells). The growth inhibition was reversible, since the growth rate recovered when the medium was changed to an RA-free one (data not shown).

Inhibition of cell infiltration into the gel by RA

As shown in Fig. 1C and D, melanoma cells rapidly infiltrated into the collagen gel and this process progressed in a time-dependent manner. The degree of

cell infiltration into the gel was remarkably suppressed as early as day 1 of culturing in both lines ($p < 0.01$).

Dose-dependency of the suppression of cell infiltration by RA

The effects of various concentrations of RA on the infiltration is shown in Table I. The effects were found to be dose-dependent in both lines. At concentrations of 10^{-6} and 10^{-5} M, the suppression rate of infiltration of S91 cells was lower than that of B16 cells ($p < 0.01$).

DISCUSSION

In the present study, rapid infiltration into the gel was observed in both melanoma cell lines (Fig. 1C, D). Because the growth rates of melanoma cells in the collagen gel were the same as those on the gel (11, 13), the increase in the number of cells in the gel was thought to be due mainly to cell infiltration. RA inhibited both cell growth and infiltration into the gel in both lines (Fig. 1), and the anti-infiltrating effect of RA was dose-dependent (Table I). The effects of RA were expressed from day 1 or 2 of culture.

The anti-proliferative effect of RA against melanoma cells has been reported by Lotan (14) and Lauharanta et al. (15). They have reported that the most sensitive cell line is Cloudman S91 and that the cell-specific differentiating function, melanogenicity, of this cell line is enhanced by the addition of RA. As shown in Fig. 1A, the growth of B16 cells was also inhibited by RA. However, the degree was lower than that of S91 cells (Fig. 1B). In contrast to S91 cells, the melanogenetic activity of B16 cells was suppressed by RA (data not shown) as previously reported by Hosoi et al. (17). On the other hand, the inhibitory effects on infiltration were observed in

Table I. Effects of various concentrations of RA on melanoma cell infiltration into the collagen gel, expressed as % of cells in gel (vs. control)

Line	Concentration of retinoic acid			
	0	10^{-7} M	10^{-6} M	10^{-5} M
B16 CBC7	100	42.5 \pm 0.8	31.4 \pm 0.3	21.2 \pm 0.5
S91	100	43.2 \pm 0.1	38.1 \pm 0.7	30.5 \pm 0.2

both lines, and the suppression rate of S91 cells was lower than that of B16 cells (Fig. 1 C, D and Table I).

The mechanism of infiltration into the gel remains to be clarified. Some authors have pointed out the important role of proteolytic enzymes in metastasis (18, 19). However, Schor et al. (9) and Bogenmann et al. (20) found no correlation between the collagenolytic activity of cells and the invasion or infiltration process. RA has been reported to be involved in the synthesis and secretion of fibronectin (FN) (21), which is known to have stimulatory (22) and inhibitory (23) effects on cell migration in plastic dishes. Schor et al. have also reported that FN is involved in cell migration in collagen gel (24). FN or another extracellular matrix, such as glycosaminoglycan, may be related to the process (25).

Recently, the anti-invasive effect of RA in mammary tumour cells has been reported (26, 27). The present study also shows that RA not only has anti-proliferative potential but also has an anti-infiltrating effect into collagen gel against melanoma cells.

As previously reported (3-8), a culture system of type I collagen gel has been considered to be a reconstituted dermis model. However, other components of the dermis including proteoglycans or fibroblasts, which will influence the behaviour of the melanoma cells, must be introduced to this system. Such an attempt has been in progress in our laboratory. The culture system using type I collagen gel as a substrate may be a useful and simple model for the *in vitro* study of tumour metastasis and invasion.

REFERENCES

1. Tickle C, Crawley A, Goodman M. Cell movement and the mechanism of invasiveness: A survey of the behavior of some normal and malignant cells implanted into the developing chick wing bud. *J Cell Sci* 1978; 31: 293-322.
2. Tickle C, Crawley A, Goodman M. Mechanism of invasiveness of epithelial tumours: Ultrastructure of the interactions of carcinoma cells with embryonic mesenchyme and epithelium. *J Cell Sci* 1978; 33: 133-155.
3. Elsdale T, Bard J. Collagen substrata for studies on cell behavior. *J Cell Biol* 1972; 54: 626-637.
4. Bell E, Ivarsson B, Merrill C. Production of a tissue-like structure by contraction of collagen lattices by human fibroblasts of different proliferative potential *in vitro*. *Proc Natl Acad Sci USA* 1979; 76: 1274-1278.
5. Yoshizato K, Taira T, Yamamoto N, Sasaki K. Remodeling of collagen: An *in vitro* model of connective tissue. *Biomedical Res* 1985; 6: 287-296.
6. Kono T, Tanii T, Furukawa M, Mizuno N, Kitajima J-I, Ishii M, Hamada T. Correlation between ageing and collagen gel contractility of human fibroblasts. *Acta Derm Venereol (Stockh)* 1990; 70: 241-244.
7. Kono T, Tanii T, Furukawa M, Mizuno N, Kitajima J, Ishii M, Hamada T, Yoshizato K. Cell cycle analysis of human dermal fibroblasts cultured on or in hydrated type I collagen lattices. *Arch Dermatol Res* 1990; 282: 258-262.
8. Schor SL. Cell proliferation and migration on collagen substrata *in vitro*. *J Cell Sci* 1980; 41: 159-175.
9. Schor SL, Schor AM, Allen TD, Winn B. The interaction of melanoma cells with fibroblasts and endothelial cells in three-dimensional macromolecular matrices: A model for tumour cell invasion. *Int J Cancer* 1985; 36: 93-102.
10. Kono T, Furukawa M, Tanii T, Kitajima J-I, Mizuno N, Ishii M, Hamada T. Contraction phenomenon of type I collagen gel by melanoma cells. *Acta Derm Venereol (Stockh)* 1990; 70: 185-188.
11. Kono T, Furukawa M, Tanii T, Mizuno N, Taniguchi S, Ishii M, Hamada T. Enhanced melanogenesis of murine melanoma cells cultured on or in the collagen gel. *Arch Dermatol Res* 1990; 282: 263-266.
12. Furukawa M, Kono T, Tanii T, Taniguchi S, Ishii M, Hamada T. Altered response to dacarbazine by murine melanoma cells cultured in type I collagen gel. *Arch Dermatol Res* 1990; 282: 278-279.
13. Furukawa M, Kono T, Tanii T, Ishii M, Hamada T, Shibata T. Proliferative potential of murine melanoma cells cultured in or on collagen gel. *J Dermatol (Tokyo)* 1990; 17: 297-302.
14. Lotan R. Effects of vitamin A and its analogs (retinoids) on normal and neoplastic cells. *Biochim Biophys Acta* 1980; 605: 33-91.
15. Lauharanta J, Kaepyahto K, Kanerva L. Changes in three-dimensional structure of cultured S91 mouse melanoma cells associated with growth inhibition and induction of melanogenesis by retinoids. *Arch Dermatol Res* 1985; 277: 147-150.
16. Oikawa A, Nakayasu M, Claunch C, Tchenn TT. Two types of melanogenesis in monolayer cultures of melanoma cells. *Cell Differ* 1972; 1: 149-155.
17. Hosoi J, Abe E, Suda T, Kuroki T. Regulation of melanin synthesis of B16 mouse melanoma cells by 1 alpha, 25-dihydroxy vitamin D₃ and retinoic acid. *Cancer Res* 1985; 45: 1474-1478.
18. Sloane BF, Honn KV, Sadler JG, Turner WA, Kimpson JJ, Taylor JD. Cathepsin B activity in B16 melanoma cells: A possible marker for metastatic potential. *Cancer Res* 1982; 42: 980-986.
19. Tryggvason K, Hoeyhtyae M, Salo T. Proteolytic degradation of extracellular matrix in tumour invasion. *Biochim Biophys Acta* 1987; 907: 191-217.
20. Bogenmann E, Mark C, Isaacs H, Neustein HB, De Clerk YA, Laug WE, Jones RA. Invasive properties of primary pediatric neoplasms *in vitro*. *Cancer Res* 1983; 43: 1176-1186.
21. Bolmer SD, Wolf G. Retinoids and phorbol esters alter release of fibronectin from enucleated cells. *Proc Natl Acad Sci USA* 1982; 79: 6541-6545.
22. Yamada KM, Olden K, Pastan I. Transformation-sensitive cell surface protein: Isolation, characterization

- and role in cellular morphology and adhesion. *Ann NY Acad Sci* 1978; 312: 256-277.
23. Couchman JR, Rees DA. The behaviour of fibroblasts migrating from chick heart explants: Changes in adhesion, locomotion and growth, and in the distribution of actomyosin and fibronectin. *J Cell Sci* 1979; 39: 149-165.
 24. Schor SL, Schor AM, Bazill GW. The effects of fibronectin on the migration of human foreskin fibroblasts and Syrian hamster melanoma cells into three-dimensional gels of native collagen fibres. *J Cell Sci* 1981; 48: 301-314.
 25. Robertson NP, Starkey JR, Hamner S, Meadows GG. Tumor cell invasion of three-dimensional matrices of defined composition: Evidence for a specific role for heparan sulfate in rodent cell lines. *Cancer Res* 1989; 49: 1816-1823.
 26. Fraker LD, Halter SA, Forbes JT. Growth inhibition by retinol of a human breast carcinoma cell line *in vitro* and athymic mice. *Cancer Res* 1984; 44: 5757-5763.
 27. Nakajima M, Lotan D, Baig MM, Carralero RM, Wood WR, Hendrix MJC, Lotan R. Inhibition by retinoic acid of type IV collagenolysis and invasion through reconstituted basement membrane by metastatic rat mammary adenocarcinoma cells. *Cancer Res* 1989; 49: 1698-1706.