

Diagnostic Value of T-SPOT.TB Test in Cutaneous Mycobacterial Infections

Yanqing CHEN^{1,2*}, Haiqin JIANG^{1,2*}, Wenyue ZHANG^{1,2}, Zhiming CHEN^{1,2}, Youming MEI^{1,2}, Hao CHEN^{1,2}, Ying SHI^{1,2}, Wei GAO^{1,2}, Le WANG^{1,2}, Santosh CHOKKAKULA^{1,2} and Hongsheng WANG^{1,2*}

¹Institute of Dermatology, Chinese Academy of Medical Sciences and Peking Union Medical College, 12 Jiangwangmiao St., Nanjing 210042, Jiangsu, and ²Jiangsu Key Laboratory of Molecular Biology for Skin Diseases and STIs, St. Nanjing, Jiangsu, China. *E-mail: whs33@vip.sina.com

*These authors contributed equally.

Accepted Jul 27, 2018; Epub ahead of print Aug 7, 2018

Detection of pathogens in skin infections caused by mycobacteria is difficult. Interferon-gamma release assay (IGRA), a preferred method for the diagnosis of latent tuberculosis infection (LTBI), has shown the capacity to detect mycobacterial infections (1–4). IGRAs have higher specificity over the tuberculin skin test, as they are based on *Mycobacterium tuberculosis* antigens encoded on RD1 (secretory antigen target 6 and culture filtrate protein 10). RD1 is absent in *Bacillus Calmette–Guérin* (BCG) and most environmental mycobacteria. However, a high degree of similarity was found in both the amino-acid sequences and the gene sequences encoding the RD1 antigens of *M. tuberculosis* and various other mycobacteria, i.e. *M. marinum*, *M. szulgai*, *M. kansasii*, and *M. goodii* (1, 5–7).

The aim of this study was to prospectively explore the diagnostic value of the T-SPOT.TB test (Oxford Immunotec Ltd., Abingdon, UK), a representative IGRA, in cutaneous infections with RD1-possessing mycobacteria, including *M. tuberculosis* and RD1-possessing nontuberculous mycobacteria (NTM). Furthermore, we evaluated the changes in T-SPOT.TB test results after antibiotic treatment.

METHODS

Between July 1, 2015 and December 30, 2017, we conducted the study at the Institute of Dermatology, Chinese Academy of Medical Sciences. All clinically suspected cases of cutaneous tuberculosis or NTM infections were screened prospectively. Ethics approval (2012-KY/LC-021) was obtained from our ethics committees. Informed consent was obtained from the participants.

There is currently no diagnostic method regarded as the gold standard for diagnosis of cutaneous tuberculosis, cutaneous infection of NTM and cutaneous fungal infection, so all eligible subjects were diagnosed with these diseases based on the following criteria: 1) chronic skin lesions, 2) pathology showing infectious granulomas, 3) molecular biological methods for the culture organism or biopsy samples confirming the corresponding disease.

Patients who met the following conditions were excluded: 1) having active tuberculosis of any organ except the skin; 2) having a history of anti-mycobacterium antibiotic treatment; 3) being ruled out by pathology; 4) refusing to participate.

Peripheral venous blood was collected for T-SPOT.TB test before treatment. Subjects in the group of cutaneous infection of RD1-possessing mycobacteria were followed up on every two months, and after 6 months of antibiotic therapy, they were further reevaluated using T-SPOT.TB test. Treatment of cutaneous tuberculosis followed the recommended guidelines by the WHO, which consist of an intensive phase for 2 months (isoniazid, rifampicin, ethambutol, and pyrazinamide) followed by a maintenance phase for 4 months (isoniazid and rifampicin) (8). Treatment for cutaneous NTM infection was designed based on drug susceptibility

tests. For those who did not participate in such tests, we utilized empiric antibiotic therapies.

The procedure of T-SPOT.TB followed the manufacturer's guidelines. The high spot numbers of Panel A minus Nil Control (PA) and Panel B minus Nil Control (PB) were recorded as spot forming cells (SFCs). Molecular biological methods were based on PCR and sequencing of the 16S rRNA and hsp65 genes (9). Homology analyses were conducted to define the strain.

Continuous variables were summarized using mean (standard deviation) for normally distributed data and using median (interquartile range (IQR)) for non-normally distributed data. The difference in age between groups was assessed by Mann–Whitney test. The differences in sex, underlying conditions, trauma history and HIV status were assessed by Chi-squared test. Positivity rates across groups were compared using Chi-squared tests. The median SFCs among groups were compared by Kruskal–Wallis test. Changes in positivity rates and SFCs before and after therapy were determined by McNemar's test and Wilcoxon rank sum test, respectively. Statistical significance was considered as $p < 0.05$. Statistical analyses were performed using IBM SPSS 19.0.

RESULTS

A total of 293 clinically suspected cases of cutaneous mycobacterial infection were screened for the study (Fig. S1¹). A total of 64 participants were eligible. Among them, 46 (71.9%) were classified as cutaneous infection of RD1-possessing mycobacterium, 8 (12.5%) as cutaneous infection of RD1-negative NTM, and 10 (15.6%) as cutaneous fungal diseases. The basic characteristics of the 64 participants are shown in Table S1¹. Forty (87.0%) of 46 cutaneous infections with RD1-possessing mycobacterium cases tested positive for T-SPOT.TB, which was much higher than those with RD1-negative NTM and fungal infections (Both 0%) ($p = 0.000$). The sensitivity in cutaneous tuberculosis and cutaneous infections of RD1-possessing NTM were 95.8% and 72.7%, respectively ($p = 0.090$) (Table S2¹).

The diagnostic performance of the T-SPOT.TB test in cutaneous infections of RD1-possessing mycobacteria is shown in Table I. Median SFCs in cutaneous infection of *M. tuberculosis*, RD1-possessing NTM, and RD1-negative NTM before anti-mycobacterium therapy were 23.5 (IQR 10.5–54.0), 14.0 (IQR 6.5–33.8) and 0.0 (IQR 0.0–1.5), respectively (Fig. S2¹) ($p = 0.000$).

All patients responded well. In the 42 patients (91.3%) who finished the study, the positive rate decreased from

¹<https://www.medicaljournals.se/acta/content/abstract/10.2340/00015555-3011>

Table I. Diagnostic performance of T-SPOT.TB in cutaneous infections of RD1-possessing mycobacterium

	Value	95% CI
Sensitivity	87.0%	73.0–94.6%
Specificity	100.0%	78.1–100.0%
Positive predictive value	100.0%	89.1–100%
Negative predictive value	75.0%	52.9–89.4%
Positive likelihood ratio	Infinity	/
Negative likelihood ratio	0.130	0.062–0.275

95.5% to 59.1% ($p=0.008$) in cutaneous tuberculosis and from 75.0% to 45.0% in RD1-possessing NTM ($p=0.031$) after therapy (Table SIII¹). Quantitative results of T-SPOT.TB in the 42 patients showed that most cases had reduction of PA or PB after treatment (Fig. S3¹). The median number of SFCs also fell significantly after treatment (Table SIII¹).

DISCUSSION

Previous reports have suggested that IGRAs show positive results for infection by NTMs such as *M. marinum*, *M. kansasii*, and *M. goodii* (1–4). In the present study, we found that cutaneous tuberculosis and RD1-possessing NTM infections also lead to positive T-SPOT.TB test results. Further, 87% of cutaneous infections by RD1-possessing mycobacteria but 0% of those by RD1-negative mycobacteria tested positive using the T-SPOT.TB test, which could potentially be used to differentiate these infections. Cutaneous tuberculosis and *M. marinum* infection are the two most common cutaneous mycobacterial diseases in China. Notably, a positive T-SPOT.TB result is a reliable indicator of the above infections, but a negative result does not rule them out.

Previous studies on the role of T-SPOT.TB in monitoring therapy have yielded varying results. We believe that this was due to significant heterogeneity between populations, methods, and study designs. All eligible cases in our study were microbiologically confirmed, thus minimizing bias brought about by uncertain cases. Despite a lack of comparability between different individuals, we found that a majority of patient SFC levels decreased after treatment, thus potentially providing a way to monitor therapeutic efficacy.

LTBI is usually diagnosed using IGRAs when no clinical or microbiological evidence of active tuberculosis can be found (10). However, we suggest that not only cutaneous tuberculosis but also RD1-possessing NTM infections should be excluded before using T-SPOT.TB test for the diagnosis of LTBI, particularly in areas with high prevalence of cutaneous infections of RD1-possessing mycobacteria. For example, areas along the river or sea, such as those found in Eastern Asia, have high numbers of skin infections caused by *M. marinum* (11–13). A number of such infections can be latent or transient while still testing positive under the T-SPOT.TB test, thus causing false-positives.

The present study certainly has limitations. The first is the limited size of the sample. This is due to limitations in current diagnostic methods, which make the diagnosis of cutaneous mycobacterial infections difficult. The second limitation is that cases of *M. kansasii*, *M. goodii*, and *M. szulgai* infection were low in number, so use of T-SPOT.TB tests for the detection of all RD1-possessing NTM infections requires further study.

ACKNOWLEDGMENTS

We thank Lemuel Shui-Lun Tsang for editing the manuscript. We thank the colleagues at the Institute of Dermatology, Chinese Academy of Medical Sciences and Peking Union Medical College for their work in sample processing.

This work was supported by grants from CAMS Innovation Fund for Medical Sciences [grant number 2016-I2M-1-005, 2017-I2M-B&R-14], Natural Science Foundation of China [grant number 81371751], and Jiangsu Provincial Natural Science Foundation [grant number BK20141065].

The authors have no conflicts of interest to declare.

REFERENCES

- Vordermeier HM, Brown J, Cockle PJ, Franken WP, Drijfhout JW, Arend SM, et al. Assessment of cross-reactivity between Mycobacterium bovis and M. kansasii ESAT-6 and CFP-10 at the T-cell epitope level. Clin Vaccine Immunol 2007; 14: 1203–1209.
- Wang MS, Wang JL, Wang XF. The performance of interferon-gamma release assay in nontuberculous mycobacterial diseases: a retrospective study in China. BMC Pulm Med 2016; 16: 163.
- Chen Y, Jiang J, Jiang H, Chen J, Wang X, Liu W, et al. Mycobacterium goodii in Patient with Facial Ulcers, Nosebleeds, and Positive T-SPOT.TB Test, China. Emerg Infect Dis 2017; 23: 1204–1206.
- Kuznetsova TI, Sauty A, Herbot CP. Uveitis with occult choroiditis due to Mycobacterium kansasii: limitations of interferon-gamma release assay (IGRA) tests (case report and mini-review on ocular non-tuberculous mycobacteria and IGRA cross-reactivity). Int Ophthalmol 2012; 32: 499–506.
- van Ingen J, de Zwaan R, Dekhuijzen R, Boeree M, van Soolingen D. Region of difference 1 in nontuberculous Mycobacterium species adds a phylogenetic and taxonomical character. J Bacteriol 2009; 191: 5865–5867.
- Andersen P, Munk ME, Pollock JM, Doherty TM. Specific immune-based diagnosis of tuberculosis. Lancet 2000; 356: 1099–1104.
- Arend SM, de Haas P, Leyten E, Rosenkrands I, Rigouts L, Andersen P, et al. ESAT-6 and CFP-10 in clinical versus environmental isolates of Mycobacterium kansasii. J Infect Dis 2005; 191: 1301–1310.
- van Zyl L, du Plessis J, Viljoen J. Cutaneous tuberculosis overview and current treatment regimens. Tuberculosis (Edinburgh, Scotland) 2015; 95: 629–638.
- Zhang X, Liu W, Liu W, Jiang H, Zong W, Zhang G, et al. Cutaneous infections caused by rapidly growing mycobacteria: case reports and review of clinical and laboratory aspects. Acta Derm Venereol 2015; 95: 985–989.
- Zumla A, George A, Sharma V, Herbert RH, Oxley A, Oliver M. The WHO 2014 global tuberculosis report – further to go. Lancet Glob Health 2015; 3: e10–12.
- Aubry A, Mougari F, Reibel F, Cambau E. Mycobacterium marinum. Microbiol Spectr 2017; 5.
- Wu J, Zhang Y, Li J, Lin S, Wang L, Jiang Y, et al. Increase in non-tuberculous mycobacteria isolated in Shanghai, China: results from a population-based study. PLoS One 2014; 9: e109736.
- Feng Y, Xu H, Wang H, Zhang C, Zong W, Wu Q. Outbreak of a cutaneous Mycobacterium marinum infection in Jiangsu Haian, China. Diagn Microbiol Infect Dis 2011; 71: 267–272.