

Phenotypic Variability with *SLURP1* Mutations and Diffuse Palmoplantar Keratoderma*

Liisa HARJAMA¹, Kaisa KETTUNEN^{2,3}, Outi ELOMAA⁴, Elisabet EINARSDOTTIR⁴⁻⁶, Hannele HEIKKILÄ¹, Sirpa KIVIRIKKO⁷, Katriina LAPPALAINEN¹, Janna SAARELA^{2,8}, Caroline ALBY⁹, Annamari RANKI¹, Juha KERE^{4,5}, Smail HADJ-RABIA¹⁰ and Katariina HANNULA-JOUUPPI^{1,4}

¹Department of Dermatology and Allergology, ERN-Skin, ²Laboratory of Genetics and ⁷Department of Clinical Genetics and Department of Medical and Clinical Genetics, University of Helsinki and Helsinki University Central Hospital, ³Institute for Molecular Medicine Finland (FIMM), Helsinki Institute of Life Science (HiLIFE), University of Helsinki, ⁴Folkhälsan Research Center, Helsinki, Finland and Research Programs Unit, Stem Cells and Metabolism Research Program, University of Helsinki, Helsinki, Finland, ⁵Department of Biosciences and Nutrition, Karolinska Institutet, Huddinge, ⁶Science for Life Laboratory, Department of Gene Technology, KTH-Royal Institute of Technology, Solna, Sweden, ⁸Norwegian Centre for Molecular Medicine (NCMM), University of Oslo, Oslo, Norway, ⁹Department of Genetics, Université Paris Descartes - Sorbonne Paris Cité, and ¹⁰Department of Dermatology and Reference Center for Genodermatoses and Rare Skin Diseases (MAGEC), FIMARAD, ERN-Skin, Université de Paris APHP5, INSERM U1163, Institut Imagine, Hôpital Universitaire Necker-Enfants Malades, Paris, France. E-mail: liisa.harjama@hus.fi

*An abstract of this manuscript was published as a poster at the 19th Annual Meeting of the European Society of Pediatric Dermatology, May, 2019. Accepted Jan 9, 2020; Epub ahead of print Jan 16, 2020

Mutations in the Secreted LY6/urokinase-type plasminogen activator receptor (uPAR)-related protein 1 gene (*SLURP1*) cause the recessively inherited palmoplantar keratodermas (PPK) Mal de Meleda (MDM, MIM 248300) and Gamborg-Nielsen (GN, MIM 244850 (1–5)). MDM is characterized by diffuse progressive and transgradient erythematous PPK, typically starting in infancy. Hyperkeratotic plaques on the elbows and knees, nail dystrophy, perioral erythema, brachydactyly, and conical-shaped fingers are reported, as well as, rarely, constriction rings with spontaneous auto-amputation of digits. Hyperhidrosis and bacterial or fungal superinfection can result in malodorous macerations, and reduced mobility of the hands and feet is common (6, 7).

GN is milder than MDM, characterized by a diffuse and transgradient hyperkeratosis with an erythematous border, and sometimes with tapered fingers (1–3, 5). The nails are normal. Distal hyperkeratosis is present only on the knuckle pads (5).

Most MDM patients are homozygous for c.82del p.(Cys28Alafs*5), c.43T>C p.(Trp15Arg) and c.286C>T p.(Arg96*) (4, 7–9), while GN patients are typically homozygous for a missense mutation c.43T>C p.(Trp15Arg)

(5). The c.82del p.(Cys28Alafs*5) variant was reported in Croatian, Tunisian and Algerian families and in a Scottish patient, suggesting a founder effect (9). This paper extends the *SLURP1* spectrum by describing phenotypic variation in 3 unrelated patients with diffuse PPK, and reporting 2 new *SLURP1* mutations.

CASE REPORTS

Case 1. An 8-year-old Finnish boy presented with diffuse waxy yellow PPK with a slight erythematous border since infancy (Table S1¹). Progressive hyperkeratosis was noted on the dorsal surface of his fingers and toes, and on his knees and elbows (Fig. 1 and Fig. S1¹). Minor wounds left hyperkeratotic scars, and water exposure turned the hyperkeratosis white and spongy. All nails were brittle with longitudinal ridges, and his teeth had enamel defects. Whole exome sequencing (Appendix S1¹) revealed a maternally inherited missense mutation, c.178G>A p.(Glu60Lys) (rs200727790) in exon 2 and a paternally inherited in-frame deletion c.218_220del p.(Cys73del) (rs748879163) in exon 3 of *SLURP1* (Fig. S2¹).

The p.(Glu60Lys) variant was predicted probably damaging by Polyphen and tolerated by SIFT (10). A total of 29 heterozygous p.(Glu60Lys) carriers, yielding a worldwide population allele frequency of 0.0001037, were reported in the GnomAD popula-

tion allele frequency database (<https://gnomad.broadinstitute.org>; last accessed 20 August 2019). There were 20 Finnish carriers (allele frequency 0.0008481). No homozygotes were reported. Only 1 Finnish heterozygous carrier for p.(Cys73del) was reported (GnomAD Finnish population allele frequency 0.000005250, worldwide population allele frequency 0.000004117).

Case 2. A 44-year-old Finnish woman had diffuse, non-progressive PPK since infancy with a waxy yellow hyperkeratosis with an erythematous border (Table S1¹). The hyperkeratosis expanded on the dorsal surface of her hands and feet, and a hyperkeratotic plaque was also present on the elbow (Fig. 1 and Fig. S1¹). She had hyperhidrosis of the hands, feet and armpits and frequent dermatophyte infections



Fig. 1. Patient phenotypes. (a–c) phenotype of case 1 c.178G>A p.(Glu60Lys) / c.218_220 p.(Cys73del), (d–f) phenotype of case 2 c.178G>A p.(Glu60Lys), and (g, h) phenotype of case 3 c.82del p.(Cys28Alafs*5) / c.286C>T p.(Arg96*).

on the affected skin. All nails were clubbed and hyperkeratotic. On water exposure the affected skin turned white and spongy. A PPK gene panel (Appendix S1¹) revealed homozygosity for *SLURP1* c.178G>A p.(Glu60Lys) (Fig. S2¹). Haplotype analysis revealed that case 1, his mother, and case 2 shared a haplotype of approximately 3,000 kb (chr8:142544410-145535881 (hg37)) surrounding the c.178G>A p.(Glu60Lys) variant, suggesting common ancestry.

Case 3. An 11-year-old Romanian boy presented with a waxy diffuse PPK with a slight erythematous border without transgrediens and progrediens since the age of 2 years (Fig. 1, Table S1¹). There was no nail, hair or tooth involvement. A custom gene panel revealed compound heterozygous mutations for a paternally inherited *SLURP1* frameshift deletion in exon 2, c.82del p.(Cys28Alafs*5) (rs587776601) and a maternally inherited truncating mutation in exon 3 c.286C>T p.(Arg96*) (rs121908317). The former causes a premature stop codon at amino acid position 32 and the latter results in the deletion of cysteine involved in the highly conserved disulphide bridges (4). Both have previously been reported homozygous in Algerian and Croatian MDM patients (4). Skin biopsies of all 3 cases revealed prominent hyperkeratosis with a very compact cornified layer, epidermal acanthosis with a pronounced granular layer and elongation of the rete ridges (Fig. S3¹). No conclusive immunohistochemistry was possible for *SLURP1*.

DISCUSSION

We report 3 PPK patients with *SLURP1* mutations and a phenotype sharing aspects of MDM and GN. Their PPK is characterized by a waxy moderately thick diffuse hyperkeratosis with an erythematous border similar to GN, although cases 1 and 2 have elbow and knee hyperkeratosis with nail abnormalities, which are rarely seen in GN, but rather in MDM. Otherwise, the manifestations are distinctly less severe than MDM, lacking, e.g. erythematous inflammation and hyperkeratosis extending widely beyond palms and soles, pseudoainhum, reduction of motility and perioral erythema. Case 3 has the mildest phenotype, with only palmoplantar hyperkeratosis. A novel feature of cases 1 and 2 is the aquagenic whitening of PPK not previously reported with MDM or GN. Our cases share similarities with milder diffuse PPK phenotypes described previously with *SLURP1* mutations c.43T>C p.(Trp15Arg) and c.256G>A p.(Gly86Arg) (9, 11, 12). We identified 2 new *SLURP1* PPK-associated variants c.178G>A p.(Glu60Lys) and c.218_220del p.(Cys73del), which are located in the central loop II region of SLURP1 (10). In SLURP2 the corresponding loop is critical in a complex formation with its receptors (13). The c.178G>A p.(Glu60Lys) is in the vicinity of the Tyr61–Pro62 peptide bond responsible for the conformational exchange in SLURP1 structure (10). The deleted Cys73 forms a disulphide bond with Cys43 in loop I of SLURP1, which may stabilize the antiparallel β -sheet structures in SLURP1 (10, 14). An 8-fold higher allele frequency of the c.178G>A variant in the Finns indicates an enrichment in Finland and the shared ~3,000 kb haplotype in its carriers suggest a Finnish founder effect.

In conclusion, we further expand the spectrum of *SLURP1* mutations with 2 novel mutations c.218_220del p.(Cys73del) and c.178G>A p.(Glu60Lys), the latter being a plausible Finnish founder mutation. Most importantly, the phenotypes of the cases reported here share features of both MDM and GN, thus verifying that these 2 can be seen as a single entity with large phenotypic variability. This suggests that *SLURP1* might be involved in PPK mimicking *KRT1* or *KRT9* mutations.

ACKNOWLEDGEMENTS

The authors acknowledge support from the Finnish Dermatological Society's (SILY) grant and Helsinki University Hospital Research Funds.

The authors have no conflicts of interest to declare.

REFERENCES

- Gamborg Nielsen P. Hereditary palmoplantar keratoderma in the northernmost county of Sweden. *Acta Derm Venereol* 1985; 65: 224–229.
- Kastl I, Anton-Lamprecht I, Gamborg Nielsen P. Hereditary palmoplantar keratosis of the Gamborg Nielsen type. Clinical and ultrastructural characteristics of a new type of autosomal recessive palmoplantar keratosis. *Arch Dermatol Res* 1990; 282: 363–370.
- Gamborg Nielsen P. Two different clinical and genetic forms of hereditary palmoplantar keratoderma in the northernmost county of Sweden. *Clin Genet* 1985; 28: 361–366.
- Fischer J, Bouadjar B, Heilig R, Huber M, Lefèvre C, Jobard F, et al. Mutations in the gene encoding SLURP-1 in Mal de Meleda. *Hum Mol Genet* 2001; 10: 875–880.
- Zhao L, Vahlquist A, Virtanen M, Wennerstrand L, Lind LK, Lundström A, et al. Palmoplantar keratoderma of the Gamborg-Nielsen type is caused by mutations in the SLURP1 gene and represents a variant of Mal de Meleda. *Acta Derm Venereol* 2014; 94: 707–710.
- Fischer J, Bouadjar B, Heilig R, Fizames C, Prud'homme JF, Weisenbach J. Genetic linkage of Meleda disease to chromosome 8qter. *Eur J Hum Genet* 1998; 6: 542–547.
- Eckl KM, Stevens HP, Lestringant GG, Westenberger-Treumann M, Traupe H, Hinz B, et al. Mal de Meleda (MDM) caused by mutations in the gene for SLURP-1 in patients from Germany, Turkey, Palestine, and the United Arab Emirates. *Hum Genet* 2003; 112: 50–56.
- Radiono S, Pramono ZAD, Oh GGK, Surana U, Widiyanti S, Danarti R. Identification of novel homozygous SLURP1 mutation in a Japanese family with Mal de Meleda. *Int J Dermatol* 2017; 56: 1161–1168.
- Nellen RG, Steijlen PM, Hennies HC, Fischer J, Munro CS, Jonkman MF, et al. Haplotype analysis in western European patients with mal de Meleda: founder effect for the W15R mutation in the SLURP1 gene. *Br J Dermatol* 2013; 168: 1372–1374.
- Shulepko MA, Lyukmanova EN, Paramonov AS, Lobas AA, Shenkarev ZO, Kasheverov IE, et al. Human neuromodulator SLURP-1: bacterial expression, binding to muscle-type nicotinic acetylcholine receptor, secondary structure, and conformational heterogeneity in solution. *Biochemistry (Mosc)* 2013; 78: 204–211.
- Zhang J, Cheng R, Ni C, Liang J, Li M, Yao Z. First Mal de Meleda report in Chinese mainland: two families with a recurrent homozygous missense mutation in SLURP-1. *J Eur Acad Dermatol Venereol* 2016; 30: 871–873.
- Bchetnia M, Bozgia M, Laroussi N, Ben Brick AS, Charfeddine C, Ben Halim N, et al. The first Mal de Meleda case in Libya: identification of a SLURP1 mutation. *Int J Dermatol* 2015; 54: 1426–1428.
- Lyukmanova EN, Shulepko MA, Kudryavtsev D, Bychkov ML, Kubaltskii DS, Kasheverov IE, et al. Human secreted Ly-6/uPAR related protein-1 (SLURP-1) is a selective allosteric antagonist of alpha7 nicotinic acetylcholine receptor. *PLoS One* 2016; 11: e0149733.
- Adeyo O, Oberer M, Ploug M, Fong LG, Young SG, Beigneux AP. Heterogeneity in the properties of mutant secreted lymphocyte antigen 6/urokinase receptor-related protein 1 (SLURP1) in Mal de Meleda. *Br J Dermatol* 2015; 173: 1066–1069.

¹<https://doi.org/10.2340/00015555-3404>