

Supplementary material to article by Enthoven P. et al. "Do pain characteristics guide selection for multimodal pain rehabilitation?"

Table SI. Patient characteristics for all unique (2011–2012) patients and for the selection employed/student/jobseeker. Values for men and women are reported separately

	All patients			Selection employed/student/jobseeker		
	All	Men	Women	All	Men	Women
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Sex	1,132 (100)	304 (26.9)	828 (73.1)	955 (100)	256 (27.1)	696 (72.9)
Country of origin						
Sweden	965 (85.2)	250 (82.2)	715 (86.4)	814 (85.2)	209 (80.7)	605 (86.9)
Europe	67 (5.9)	20 (6.6)	47 (5.7)	54 (5.7)	19 (7.3)	35 (5.0)
Non-European	99 (8.7)	34 (11.3)	65 (7.9)	86 (9.0)	31 (12.0)	55 (7.9)
Missing	1 (0.1)	0 (0.0)	1 (0.1)	1 (0.1)	0 (0.0)	1 (0.1)
Education						
University/College	213 (18.8)	45 (14.8)	168 (20.3)	194 (20.3)	42 (16.2)	152 (21.8)
High school	630 (55.7)	174 (57.2)	456 (55.1)	546 (57.2)	148 (57.1)	398 (57.2)
Compulsory	215 (19.0)	63 (20.7)	152 (18.4)	155 (16.2)	50 (19.3)	105 (15.1)
Missing (=övrigt)	74 (6.5)	22 (7.2)	52 (6.3)	60 (6.3)	19 (7.3)	41 (5.9)
Working status						
Employed	627 (55.4)	181 (59.5)	446 (53.9)	627 (65.7)	181 (59.5)	446 (53.9)
Student	82 (7.2)	10 (3.3)	72 (8.7)	82 (8.6)	10 (3.3)	72 (8.7)
Jobseeker	246 (21.7)	68 (22.4)	178 (21.7)	246 (25.8)	68 (22.4)	178 (21.7)
Not gainfully employed	117 (15.6)	45 (14.8)	132 (15.6)			
Pain location variation						
Pain location constant	770 (68.0)	226 (74.3)	544 (65.7)	653 (68.4)	192 (74.1)	461 (66.2)
Pain location varies	362 (32.0)	78 (25.7)	284 (34.3)	302 (31.6)	67 (25.9)	235 (33.8)
Mental health						
Anxiety (> 10)	392 (34.6)	111 (36.5)	281 (33.9)	316 (33.1)	94 (36.3)	222 (31.9)
Missing	18 (1.6)	9 (3.0)	9 (1.1)	14 (1.5)	9 (3.5)	5 (0.7)
Depression (> 10)	342 (30.2)	108 (35.5)	234 (28.3)	263 (27.5)	85 (32.8)	178 (22.6)
Missing	14 (1.2)	6 (2.0)	8 (1.0)	11 (1.2)	6 (2.3)	5 (0.7)
Selected for MMR	304 (26.9)	43 (14.1)	253 (30.6)	279 (29.2)	41 (15.8)	238 (34.2)
	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)
Age	41.1 (12.1)	44.3 (11.6)	40.0 (12.1)	40.2 (11.6)	43.2 (11.2)	39.2 (11.6)
Pain intensity in the previous week (NRS)	7.0 (1.8)	6.8 (1.9)	7.1 (1.7)	6.9 (1.8)	6.7 (1.9)	7.0 (1.7)
Pain severity (MPI)	4.5 (0.9)	4.4 (1.0)	4.5 (0.9)	4.5 (0.9)	4.4 (1.0)	4.5 (0.9)
Pain interference with daily life (MPI)	4.5 (1.1)	4.5 (1.1)	4.5 (1.0)	4.4 (1.1)	4.5 (1.2)	4.4 (1.0)
Number of days since pain initiated	2,964 (3,066)	2,955 (3,528)	2,967 (2,871)	2,700 (2,887)	2,652 (3,405)	2,718 (2,664)
Number of days with chronic pain	2,335 (2,658)	2,437 (3,297)	2,295 (2,366)	2,188 (2,597)	2,211 (3,272)	2,179 (2,293)
Number of pain sites	13.8 (8.3)	11.6 (7.6)	14.6 (8.4)	13.5 (8.2)	11.6 (7.7)	14.3 (8.3)
Number of quadrants with pain	3.3 (1.1)	3.2 (1.1)	3.3 (1.0)	3.2 (1.1)	3.2 (1.1)	3.3 (1.1)
1PCS SF36	28.9 (8.3)	29.3 (8.6)	28.8 (8.1)	29.4 (8.2)	29.4 (8.7)	29.4 (8.0)
MCS SF36	36.2 (12.8)	35.4 (12.9)	36.5 (12.8)	37.9 (12.8)	36.1 (13.0)	37.2 (12.7)
EQ-5D Index	0.27 (0.31)	0.23 (0.32)	0.28 (0.31)	0.29 (0.31)	0.25 (0.33)	0.31 (0.31)
EQ-5D VAS	38.1 (20.8)	37.6 (20.6)	38.2 (20.9)	38.9 (20.7)	38.8 (20.9)	40.0 (20.6)
Tampa Scale	39.7 (9.1)	42.6 (10.0)	38.5 (8.5)	39.3 (9.1)	42.5 (10.2)	37.9 (8.3)

Unique patients: patients who are not included in the study by Haukenes et al. 2015.

MMR: multimodal rehabilitation; NRS: numeric rating scale; MPI: Multidimensional Pain Inventory; PCS SF36: Physical Component Summary Short Form 36; MCS: mental Component Summary; EQ-5D: European Quality of Life Instrument; Tampa Scale: Tampa Scale of Kinesiophobia.