

Supplementary material to article by G. Francisco et al. “Adult spasticity international registry study: methodology and baseline patient, healthcare provider, and caregiver characteristics”

**Table SI.** Spasticity treatment modalities at baseline by aetiology

	Cerebral palsy (n = 77)	Multiple sclerosis (n = 119)	Spinal cord injury (n = 42)	Stroke (n = 411)	Traumatic brain injury (n = 45)	Other (n = 72)	Total (n = 730)
Acupuncture, n (%)	74 2 (2.7)	115 12 (10.4)	41 1 (2.4)	392 30 (7.7)	44 7 (15.9)	70 9 (12.9)	701 56 (8.0)
Assistive devices, n (%)	74 37 (50.0)	115 96 (83.5)	41 30 (73.2)	392 282 (71.9)	44 31 (70.5)	70 45 (64.3)	701 494 (70.5)
Casting, n (%)	74 16 (21.6)	115 2 (1.7)	41 5 (12.2)	392 42 (10.7)	44 8 (18.2)	70 7 (10.0)	701 72 (10.3)
Chemodenervation, n (%)	74 1 (1.4)	115 6 (5.2)	41 2 (4.9)	392 20 (5.1)	44 1 (2.3)	70 2 (2.9)	701 30 (4.3)
Intrathecal therapy, n (%)	74 10 (13.5)	115 11 (9.6)	41 8 (19.5)	392 37 (9.4)	44 7 (15.9)	70 9 (12.9)	701 75 (10.7)
Orthotics, n (%)	74 36 (48.6)	115 39 (33.9)	41 18 (43.9)	393 241 (61.3)	44 27 (61.4)	70 36 (51.4)	702 377 (53.7)
Physio or occupational therapy, n (%)	74 53 (71.6)	115 89 (77.4)	41 34 (82.9)	393 333 (84.7)	44 35 (79.5)	70 58 (82.9)	702 575 (81.9)
Surgeries or procedures, n (%)	74 31 (41.9)	115 2 (1.7)	41 10 (24.4)	393 37 (9.4)	44 14 (31.8)	70 13 (18.6)	702 93 (13.2)

NOTE: Allowed responses included “yes” or “not mentioned”.