

Table SIV. Disciplines involved in delivering sexual rehabilitation following stroke: priorities and consensus

| Primary clinician delivering intervention | Consensus score (IQR/2) | Interpretation | Priority based on percentage of responders identifying the clinician in their top 3 | | Priority based on mean priority scores (1–5) | |
|--|-------------------------|----------------|---|----------------|--|----------------|
| | | | Responders selecting topic in their top 3 (%) | Interpretation | Mean | Interpretation |
| Geriatrician | 1 | Moderate | 8.5 | Low | 3.1 | High |
| Neurologist | 0.8 | High | 12.8 | Low | 3.5 | High |
| Rehabilitation Physician | 0.5 | High | 42.6 | High | 3.8 | High |
| Nurse working in acute stroke or rehabilitation services | 0.5 | High | 24.5 | Intermediate | 3.8 | High |
| Nurse with specialist knowledge in stroke and sexuality | 0.5 | High | 63.8 | High | 4.4 | High |
| Occupational therapist | 1 | Moderate | 41.5 | High | 3.8 | High |
| Physiotherapist | 0 | High | 22.3 | Intermediate | 3.8 | High |
| Psychologist | 0.3 | High | 35.1 | High | 3.7 | High |
| Sexologist/sex educator/counsellor | 0.5 | High | 45.7 | High | 4.0 | High |
| Social worker | 1 | Moderate | 22.3 | Intermediate | 3.5 | High |
| Speech pathologist | 1 | Moderate | 19.1 | Intermediate | 3.4 | High |