

**Table SII.** International Classification of Functioning, Disability and Health (ICF) categories in Body Functions, showing significant differences between 2 different evaluation methods (test-based ICF assessment at the start of rehabilitation vs observation-based ICF assessment at the end of rehabilitation)

ICF categories	None 0-4% n (%)	Mild 5-24% n (%)	Moderate 25-49% n (%)	Severe 50-95% n (%)	Complete 96-100% n (%)
Immunological functions (b435)					
Test-based, M = 0	29 (52)	24 (43)	3 (5)	0 (0)	0 (0)
Observation-based, M = 1 ( $p < 0.02$ and $< 0.02$ )	14 (35)	16 (40)	6 (15)	4 (10)	0 (0)
Basic interpersonal interactions (d710)					
Test-based, M = 1 ( $p < 0.005$ and $< 0.005$ )	21 (42)	15 (30)	7 (14)	7 (8)	0 (0)
Observation-based, M = 0	30 (71)	11 (26)	1 (2)	0 (0)	0 (0)
Informal socializing (d9205)					
Test-based, M = 2 ( $p < 0.005$ and $< 0.05$ )	7 (14)	9 (19)	13 (27)	19 (40)	0 (0)
Observation-based, M = 1	4 (10)	18 (44)	16 (39)	3 (7)	0 (0)

Scores "non-specified" and "non-applicable" were not used for comparison and were considered as missing values.  $\chi^2$  and Mann-Whitney test were used for statistical analysis (presented in parentheses, respectively). M=median.