



INFLUENCE OF ARM CRANK ERGOMETRY ON DEVELOPMENT OF LYMPHOEDEMA IN BREAST CANCER PATIENTS AFTER AXILLARY DISSECTION: A RANDOMIZED CONTROLLED TRIAL

Thorsten SCHMIDT, PhD¹, Jette BERNER², Walter JONAT, MD, PhD², Burkhard WEISSER, MD, PhD³, Christoph RÖCKEN, MD, PhD¹, Marion VAN MACKELENBERGH, MD² and Christoph MUNDHENKE, MD, PhD²

From the ¹Comprehensive Cancer Center North, ²OB/GYN, University Hospital for Women, and ³Institute of Sport Science, University of Kiel, Kiel, Germany

Objective: To investigate the safety and efficacy of arm crank ergometry in breast cancer patients after axillary lymph node dissection, with regard to changes in bioelectrical impedance analysis, arm circumference, muscular strength, quality of life and fatigue.

Design: Randomized controlled clinical intervention trial.

Subjects: Forty-nine patients with breast cancer after axillary lymph node dissection.

Methods: Arm crank ergometer training twice-weekly was compared with usual care over 12 weeks.

Results: The arm crank ergometer group improved significantly in terms of lean body mass and skeletal muscle mass, and showed a significant decrease in body fat. In the arm crank ergometer group, as well as the usual care group, a significant increase in armpit circumference was detected during the training period. The magnitude of the gain was higher in the usual care group. For all other measured regions of the arm a significant decrease in circumference was seen in both groups. Muscular strength of the upper extremity increased significantly in both groups, with a greater improvement in the arm crank ergometer group. In both groups a non-significant trend towards improvement in quality of life was observed. The arm crank ergometer group showed significant improvements in physical functioning, general fatigue and physical fatigue.

Conclusion: These results confirm the feasibility of arm crank ergometer training after axillary lymph node dissection and highlight improvements in strength, quality of life and reduced arm symptoms with this training.

Key words: breast cancer-related lymphoedema; arm crank ergometer; physical activity.

Accepted Sep 14, 2016; Epub ahead of print Dec 8, 2016

J Rehabil Med 2017; 47: 78–83

Correspondence address: Thorsten Schmidt, Breast Unit, Comprehensive Cancer Center North, University of Kiel, Arnold-Heller-Str. 3, DE-24105 Kiel, Germany. E-mail: thorsten.schmidt@uksh.de

Breast cancer is the most prevalent cancer in women, with a high survival rate but significant long-term morbidity after treatment. Rehabilitation of post-treatment effects of breast cancer treatment

is needed to improve functioning and quality of life (QoL). In addition to surgery of the breast, 61% of patients undergo axillary sentinel lymph node biopsy. Thirty-three percent of patients with breast cancer require complete axillary lymph node dissection (ALND) (1, 2). The risk of developing a lymphoedema, based on the clinical definition, is 5–20% (1, 3, 4).

Rehabilitation of breast cancer patients focuses on side-effects after axillary surgery, such as pain and physical impairments. Meanwhile, there is evidence that physical activity is an important factor in improving overall survival and time to progression (5, 6). Inactivity, related or not related to cancer treatment, can weaken the skeleton, cause muscle loss and lead to fat gain (7–10). These changes in body composition place breast cancer survivors at a higher risk of obesity-related diseases, breast cancer recurrence, frailty and fractures (11). Previous findings suggest that physical activity during and after breast cancer treatment can not only reduce side-effects (8, 12–15), but also improve clinical outcome (9, 10, 12, 14–16).

In the past, intense physical activity of the upper extremity has not been recommended after ALND and/or radiation to the axilla. As there is a lack of recommendations or guidelines on physical activity for patients with lymphoedema of the arm after breast cancer therapy, women with this postoperative complication are uncertain about exercising. Recent studies have reported that progressive weight-lifting did not have any preventative effect on development of breast/arm swelling (or lymphoedema). Nonetheless, there are only a small number of randomized controlled trials on this topic (16–25).

This clinical interventional trial investigates the feasibility of arm crank ergometer (ACE) training after ALND for primary breast cancer. Safety, efficacy and the influence on prevention or improvement of existing lymphoedema of the arm are the main study endpoints.

MATERIAL AND METHODS

Design and procedures

In this prospective, controlled, randomized intervention trial 12-week supervised ACE-training was compared with usual care (UC) in breast cancer patients after ALND. Patients were

recruited at a major, academic, breast oncology unit in Germany. Participating women had to meet the following eligibility criteria: primary breast cancer; within 3 years after axillary lymph node dissection; age range 18–75 years. Exclusion criteria were: intended radiotherapy or chemotherapy during the study period; acute infectious disease; severe cardiac disease (New York Heart Association functional class III; myocardial infarction <3 months); severe pulmonary or renal insufficiency (glomerular filtration rate <30%); serious neurological disorders; fewer than 10,000 platelets/ml; and haemoglobin <8g/dl.

The study was approved by the local ethics committee (registration number: AZA 104/14). All participants were included in the study after providing written informed consent, as required by the Declaration of Helsinki (1975).

Randomization

After baseline assessments the patients were assigned randomly (1:1) to supervised ACE-training or UC using a computer-generated programme. The allocation sequence was executed by the clinical research unit and concealed from the project team. To prevent a possible bias, research physicians did not have access to the randomization files.

Statistical analysis

The analysis included data for patients who attended at least 70% of the scheduled training sessions. Performance was assessed at baseline (T1) and 12 weeks after initiation of physical intervention (T2). Primary endpoints of this trial were change in body composition of the upper extremity, measured by bioelectrical impedance analysis (BIA), arm volume (arm circumference), muscular strength (NM), QoL (European Organization for Research and Treatment of Cancer (EORTC) QLQ C30+BR23) and fatigue (Multidimensional Fatigue Inventory; MFI 20) before training and after 12 weeks of training.

For the within-group changes, a dependent *t*-test (body composition of the upper extremity and the trunk, arm volume (arm circumference) and muscular strength) and the Wilcoxon-test (EORTC and fatigue questionnaire) were used. Comparisons of the ACE group with the UC group for changes from baseline to 12-week assessment (T2–T1) were performed using independent *t*-test. Statistical significance for the *t*-test at T1 and T2 was set at a probability level of $p < 0.05$. The effects are expressed with mean and standard deviations. Analysis was performed using the SPSS system for windows (Version: PASW 21).

Setting and participants

Body mass index (BMI) was calculated and body composition was determined by anthropometry and bioelectrical impedance analysis (BIA) (InBody, Modell 520, Germany). The body composition of the upper extremity and the trunk were measured under constant conditions (e.g. time of day and good hydration). BIA determined values (in %) for lean body mass (LBM), skeletal muscle mass, and body fat. The participant was positioned in an orthostatic position on a platform with lower electrodes for the feet and upper electrodes for the hands, enabling evaluation of the impedance of each body segment.

Arm circumference was measured in all randomized patients at defined anatomical regions (armpit, 15 cm above the lateral epicondyles of the elbow, elbow, and wrist). This measurement was performed on the arm (after ALND) and, for comparison, on the contralateral arm (non-ALND arm). Measurements were performed at time-points T1 and T2 (before and after study-related training). The impact of the training type (ACE vs UC)

on changes in arm circumferences was statistically analysed. Lymphoedema was defined as a difference of the arm circumference >5% between the time of measurement.

The muscular strength of the upper extremities (latissimus dorsi muscle and pectoralis muscle) was assessed by measuring their isometric muscular capacity with M3 Diagnos (Schnell, Germany) combined with “Diagnos Professional 2000” software. To test the maximum strength of the upper extremities the patients were encouraged to stretch their arms against the fixed bar for approximately 5 s (pectoralis muscle), followed by pulling the fixed bar for approximately 5 s (latissimus dorsi muscle). Every test was repeated twice, with appropriate breaks in between.

The EORTC has developed several validated questionnaires to assess the QoL of patients with cancer in a multidimensional approach. The questionnaire used was EORTC QLQ C30, version 3 BR23, which was especially developed for patients with breast cancer. The results section of the questionnaire focuses on QoL and fatigue (26, 27). Fatigue symptoms were assessed with the Multidimensional Fatigue Inventory with 20 questions (MFI-20). The MFI-20 is an internationally frequently used questionnaire (28, 29).

The ACE training included strength and endurance training with an arm crank ergometer (MOTomed®, Reck, Betzenweiler) for 60 min twice weekly for 12 weeks. Every training session was supervised and documented by experienced exercise therapists. Before each training session, the intensity levels were checked according to the exercise guidelines for cancer survivors of the American College of Sports Medicine (30). The training principle consists of 3 possible training components: 1. step, a passive completely motorized training; 2. step, a motor-supported self-training; and 3. step, a training by patient-generated power. Depending on the patient-reported performance status, patients participated in the different training modalities. In case of a reduced performance status they enhanced their performance gradually and proceeded stepwise from training steps 1 to 3. When they reported a good performance status, they entered training steps 2 or 3.

The Borg scale was used as a subjective reference point for self-reported performance status during training. Patients were encouraged to be active at Borg levels 11–14 (31, 32). The 45-min training included a 10-min warm-up, 25–30 min of exercise (in training steps 1–3) and a 5-min cool-down period. When a lymphoedema occurred the training intensity was reduced and carried out either motorized or motor-supported. A lymphoedema was defined as a 5% increase in the volume of the affected limb. Participants in the usual care group were informed about the feasibility of physical activity and the impact of physical activity for breast cancer-related lymphoedema, but did not receive supervised training.

RESULTS

A total of 120 patients were identified retrospectively and contacted between May and December 2014. Sixty-two patients fulfilled the inclusion criteria and were enrolled and randomized into the ACE group or UC group. Fifty-eight of the identified patients were unable to participate; of these, 46 declined to participate due to the distance to the study centre, 8 had died, and 4 were not interested. Due to timing issues, family problems, or withdrawal of consent, 13 out of 62 enrolled and randomized patients dropped out: 6 withdrew from the ACE group and 7 from the UC group. Data

Table I. Patient characteristics

	ACE group	UC group
Total, <i>n</i>	21	28
Age, years, mean (SD)	61.67 (10.01)	53 (10.7)
BMI, mean (SD)	30.42 (7.55)	26.92 (6.42)
Type of breast surgery, <i>n</i>		
Modified radical mastectomy	10	12
Subcutaneous mastectomy	-	1
Breast-conserving surgery	11	15
Type of postoperative radiotherapy, <i>n</i>		
RT of the breast and thoracic wall	13	15
RT of the breast, thoracic wall and lymphatic regions	8	11
No RT	0	3
Month since completion of therapy, mean (SD)	9.66 (3.61)	10.27 (2.56)

BMI: body mass index; SD: standard deviation; UC: usual care; ACE: arm crank ergometer; RT: radiotherapy.

for 49 patients were fully evaluable (21 patients in the ACE group, 28 in the UC group). An overview of the patient cohort is shown in Table I and Fig. 1.

During the course of the study a significant increase in lean body mass (LBM) of the patients in the ACE group was detected (ALND-arm, $p=0.017$; non-ALND-arm, $p=0.004$). Furthermore, a significant increase in skeletal muscle mass (ACE group, $p=0.049$) and decrease in body fat in the ACE group (ACE group, $p=0.009$) could be observed. In the UC group there was a trend towards improvement, but no significant change, in LBM (ALND-arm, $p=0.679$; non-ALND-arm, $p=0.354$), skeletal muscle mass (UC group, $p=0.385$) or body fat (UC group, $p=0.393$) (Table II).

A decrease in circumference of the ALND-arm could be detected at the measuring point's wrist joint, 15 cm above the lateral epicondylus of the elbow and the




Fig. 1. Study flowchart. ACE: arm crank ergometer; UC: usual care.

elbow in patients in both study groups. Also, a significant increase in the circumference of the armpit of the ALND-arm could be seen in both groups, whereby the increase was stronger in the UC group (ACE group: +4.29%, UC group: 10.66%) (Table III). In comparison with the ALND-arm, the non-ALND-arm showed no major difference in both groups.

Muscular strength improved in both groups from T1 to T2. Compared with the control group, the ACE group had a greater improvement in muscular strength of the upper extremity (non-significant) (Table IV).

In both groups a non-significant trend towards

Table II. Body composition of the upper extremity of the intervention group and the usual care group at baseline (T1) vs 12 weeks after the initiation of intervention (T2)

Parameter	T1		T2		T1 vs T2 difference <i>p</i> -value ^a	
	ACE group Mean (SD)	UC group Mean (SD)	ACE group Mean (SD)	UC group Mean (SD)	ACE	UC
Lean body mass ALND-arm, kg	2.7 (0.77)	2.55 (0.53)	2.79 (0.81)	2.57 (0.52)	0.017	0.679
Lean body mass non-ALND-arm, kg	2.66 (0.73)	2.54 (0.53)	2.72 (0.76)	2.52 (0.52)	0.004	0.354
Skeletal muscle mass, kg	26.92 (6.03)	26.16 (4.21)	28.09 (6.65)	26.45 (4.26)	0.049	0.385
Body fat, kg	35.62 (17.32)	25.92 (13.66)	34.04 (17.21)	26.27 (13.59)	0.009	0.393

T1: baseline; T2: 12 weeks after initiation of the intervention, ^a*p*-values were calculated using dependent *t*-test. ACE: arm crank ergometer; UC: usual care; ALND: axillary lymph node dissection.

Table III. Circumference of the upper extremity of the intervention group and the usual care group at baseline (T1) vs 12 weeks after the initiation of intervention (T2)

Parameter	T1		T2		T1 vs T2 difference %		T1 vs T2 difference <i>p</i> -value ^a	
	ACE group Mean (SD)	UC group Mean (SD)	ACE group Mean (SD)	UC group Mean (SD)	ACE	UC	ACE	UC
Arm circumference ALND-arm wrist joint, cm	17.69 (1.65)	17.09 (1.46)	17.14 (1.65)	16.35 (1.29)	-3.1	-4.3	0.038	0.01
Arm circumference ALND-arm 15 cm above the lateral epicondylus of the elbow, cm	34.24 (5.85)	31.42 (4.47)	33.79 (5.56)	30.92 (4.22)	-1.31	-1.50	0.232	0.01
Arm circumference ALND-arm of the elbow, cm	28.26 (3.98)	26.71 (2.43)	27.6 (3.43)	26.08 (2.59)	-2.33	-2.25	0.027	0.001
Arm circumference ALND-arm of the armpit, cm	39.31 (7.2)	36.77 (4.84)	41.71 (5.19)	40.69 (4.67)	+4.29	+10.66	0.025	0.01

T1: baseline; T2: 12 weeks after the initiation of intervention, ^a*p*-values were calculated using dependent *t*-test. ACE: arm crank ergometer; UC: usual care; ALND: axillary lymph node dissection.

Table IV. Muscular strength of the upper extremity of the intervention group and the usual care group at baseline (T1) vs 12 weeks after initiation of the intervention (T2)

	T1		T2		%		<i>p</i> -value ^a	
	ACE group Mean (SD)	UC group Mean (SD)	ACE group Mean (SD)	UC group Mean (SD)	ACE group Mean (SD)	UC group Mean (SD)	ACE	UC
Muscular strength of								
Muscle pectoralis ALND-arm, NM	57.31 (34.52)	59.46 (22.67)	81.88 (54.29)	79.46 (40.62)	42.8	33.6	0.002	0.011
Muscle pectoralis control-arm, NM	61.14 (35.69)	61.46 (21.94)	87.33 (57.16)	74.54 (35.88)	42.8	21.2	0.005	0.038
Muscle latissimus doris ALND-arm, NM	63.43 (29.98)	70.88 (29.21)	86.90 (41.85)	86.92 (34.64)	37.0	22.6	0.01	0.043
Muscle latissimus doris control-arm, NM	66.43 (30.79)	71.81 (27.08)	88.5 (41.7)	87.42 (34.91)	33.2	21.7	0.01	0.034

T1: baseline; T2: 12 weeks after the initiation of intervention, ^a*p*-values were calculated using dependent *t*-test.

SD: standard deviation; UC: usual care; ALND: axillary lymph node dissection; ACE: arm crank ergometer; NM: muscular strength.

Table V. Psychology parameters, quality of life (EORTC QLQ C30) fatigue (MFI20) and total activity score (IPAG) of the intervention group and the usual care group

Parameter	T1				T2				<i>p</i> -value	
	ACE group		UC group		ACE group		UC group		ACE group	UC group
	Median	Mean (SD)	Median	Mean (SD)	Median	Mean (SD)	Median	Mean (SD)		
EORTC										
Physical functioning	66.67	63.49 (18.93)	86.67	83.1 (16.97)	66.67	68.57 (16.97)	86.67	85 (13.53)	0.039	0.428
Quality of life (global health status)	66.67	60.32 (17.46)	62.5	63.69 (24.66)	66.67	64.68 (16.22)	66.67	68 (16.21)	0.101	0.214
MFI 20										
General fatigue	12	12.81 (4.21)	10.5	10.39 (5.19)	11	11.29 (4.15)	10	9.89 (3.82)	0.032	0.483
Physical fatigue	13	13.24 (3.506)	8.5	9.86 (5.275)	10	11 (3.72)	9.5	9.43 (4.194)	0.002	0.42

T1: Baseline; T2: 12 weeks after the initiation of intervention, *p* values were calculated using Wilcoxon-test; ACE: Arm crank ergometer, UC: Usual Care; SD: standard deviation.

improvement in QoL was observed (ACE group, $p=0.101$; UC group, $p=0.202$). The ACE group significantly improved in terms of physical functioning (ACE group, $p=0.038$; UC group, $p=0.428$); general fatigue (ACE group, $p=0.032$; UC group, $p=0.483$) and physical fatigue (ACE group, $p=0.002$; UC group, $p=0.42$). In the UC group only minor non-significant changes were detected. Other scales of the EORTC and the BR23 questionnaire module showed no changes in the ACE group and UC group (Table V).

A comparison of the mean differences between the ACE group and the UC group showed no significant differences between the 2 groups. In both groups the range of the circumference of the upper extremity decreased at the measuring points wrist joint, 15 cm above the lateral epicondylus of the elbow and the elbow. In both groups an increased mean difference in the circumference of the armpit of the ALND-arm was seen (Table VI)

A similar picture emerged for the data regarding body composition and muscular strength, whereby, in comparison with the UC group, the ACE group showed a major improvement (Tables VII and VIII).

DISCUSSION

Until recently it was not recommended that patients undertake physical activity of the upper extremity after axillary lymph node dissection and/or radiation to the axilla. There was unsubstantiated concern that physical activity could initiate or lead to deterioration of a lymphoedema. A small number of publications showed that physical activity and upper-body exercise does not have an influence on the development or worsening of lymphoedema (20). The majority of publications investigated the impact of resistance training, whereas

Table VII. Mean difference in body composition of the upper extremity of the intervention group and the usual care group

Parameter	ACE group Mean (SD)	UC group Mean (SD)	<i>p</i> -value ^a
Lean body mass ALND-arm	0.086 (0.143)	0.015 (0.19)	0.182
Lean body mass non-ALND-arm	0.058 (0.078)	-0.019 (0.105)	0.009
Skeletal muscle mass	1.168 (2.417)	0.292 (1.684)	0.159
Body fat	-1.57 (2.35)	0.35 (2.05)	0.006

^a*p*-values were calculated using independent *t*-test.

SD: standard deviation; ACE: arm crank ergometer; UC: usual care; ALND: axillary lymph node dissection.

Table VI. Mean difference (T2-T1) in the circumference of the upper extremity of the intervention group and the usual care group

Parameter	ACE group Mean (SD)	UC group Mean (SD)	<i>p</i> -value ^a
Arm circumference ALND-arm wrist joint	-0.523 (0.732)	-0.480 (0.519)	0.815
Arm circumference ALND-arm 15 cm above the lateral epicondylus of the elbow	-0.595 (1.570)	-0.46 (1.41)	0.76
Arm circumference ALND-arm of the elbow	-0.667 (1.27)	-0.634 (0.889)	0.92
Arm circumference ALND-arm of the armpit	1.889 (3.26)	3.92 (4.00)	0.082

^a*p*-values were calculated using independent *t*-test.

SD: standard deviation; ACE: arm crank ergometer; UC: usual care; ALND: axillary lymph node dissection.

Table VIII. Mean difference (T2-T1) in the muscular strength of the upper extremity of the intervention group and the usual care group

Muscular strength of	ACE group	UC group	p-value ^a
Muscle pectoralis ALND-arm	24.57 (32.20)	20.00 (37.31)	0.66
Muscle pectoralis control-arm	26.19 (38.38)	13.07 (30.42)	0.198
Muscle latissimus doris ALND-arm	23.47 (18.22)	16.03 (38.28)	0.387
Muscle latissimus doris control-arm	22.07 (20.29)	15.61 (35.54)	0.439

^ap-values were calculated using independent t-test.

ACE: arm crank ergometer; UC: usual care; ALND: axillary lymph node dissection.

upper-body endurance training was under-represented. The primary objective of our study was to determine the impact of arm crank ergometer training as a strength and endurance training on the development or worsening of lymphoedema in breast cancer patients who have undergone axillary lymph node dissection. Based on the inclusion criteria comparison with the few published randomized controlled trials is difficult. Sagen et al. included women after an axillary dissection (18), Schmitz et al. (23) included women who had had a sentinel node biopsy or an axillary dissection (19). Blommquist et al. did not report any details about the type of surgery. Furthermore, different definitions for lymphoedema were used. Schmitz et al. defined a lymphoedema as an increase in the volume of the affected limb of 5% or more, accompanied by an increase of 5% or more in the difference in the volume or circumference of the affected and unaffected limbs (19). In the study by Bloomquist et al. a lymphoedema is defined as a difference of ≤ 2 cm at 2 or more measures (23).

Similar to the results of previous studies, strength and endurance training with an arm crank ergometer did not have a detrimental effect on breast cancer-associated lymphoedema compared with usual care. A decrease in arm circumference was seen in both groups and potential lymphoedema exacerbations were found in the UC group. Furthermore, muscle strength improved in both groups, whereas the improvement in the ACE group was more pronounced. A possible explanation for the decrease in arm circumference and the improvement in muscle strength in the UC group is that participants in the UC group were more confident to exercise their upper limb due to education about the impact of physical activity on breast cancer-related lymphoedema. Furthermore, physical activity is known to improve the flow of lymphatic fluid and protein resorption (20).

The improvement in QoL and physical functioning detected in the ACE group is confirmed by other study results. In a population-based cohort study Chen et al. evaluated the effect of regular exercise on QoL during the first 3 years after breast cancer diagnosis. It was observed that women with a higher exercise score were more likely to have higher scores for total QoL

and specific subscales, compared with women taking non-regular exercise after breast cancer diagnosis (33). Gautam et al. confirmed the results in a pre-post intervention study with breast cancer patients who had undergone mastectomy and had completed medical therapy. After an 8-week home-based programme the participants' QoL had improved (25). The improvement in QoL in the UC group was the same as that in the ACE group. In comparison with the literature the evidence grade is high for physical factors and low for mental health data (QoL, fatigue, anxiety, depression), as published trials differ substantially in design and the study populations and cannot be compared head to head. The current literature in the field of QoL of breast cancer patients after medical treatment confirmed our data (34, 35). The ACE group improved significantly in terms of general fatigue and physical fatigue, whereas non-significant changes could be detected in the UC. In fact, there are limited studies investigating the impact of physical activity on fatigue.

This is one of the first clinical trials to investigate the safety and efficacy of arm crank ergometry for endurance and strength training.

A limitation of this study is that the results are limited by the large amount of parallel statistical testing performed. The probability of occurrence of false-positive results was thereby increased. Furthermore, the small sample size limited the power of the results. It is therefore our intention to verify the results in a follow-up study with increased sample size and subgroup analysis.

In conclusion, the results of the present study, supported by the current literature, show a positive development of strength, endurance, QoL and fatigue and no influence of the onset or worsening of lymphoedema. These positive results suggest that strength and endurance training with an arm crank ergometer should be implemented in the regular aftercare of patients with breast cancer.

ACKNOWLEDGEMENTS

The author would like to acknowledge Wiebke Kaczmarek and Katrin Dehn of the Institute of Sport Science, Kiel, Germany for their support in patient care.

This trial was supported by Reck, Betzweiler, Germany with financial support. The Supporter had no role in the study design, data collection and analysis, preparation of the manuscript or decision to publish.

The authors declare no conflicts of interest.

REFERENCES

1. Jemal A, Siegel R, Ward E, Hao Y, Xu J, Thun MJ. Cancer statistics 2009. *CA Cancer J Clin* 2009; 59: 225-249.

2. Francis WP, Abghari P, Du W, Rymal C, Suna M, Kosir MA. Improving surgical outcomes: standardizing the reporting of incidence and severity of acute lymphedema after sentinel lymph node biopsy and axillary lymph node dissection. *Am J Surg* 2006; 192: 636–639.
3. Hayes SC, Janda M, Cornish B, Battistutta D, Newman B. Lymphedema after breast cancer: incidence, risk factors, and effect on upper body function. *J Clin Oncol* 2008; 26: 3536–3542.
4. Wilke LG, McCall LM, Posther KE, Whitworth PW, Reintgen DS, Leitch AM, et al. Surgical complications associated with sentinel lymph node biopsy: results from a prospective international cooperative group trial. *Ann Surg Oncol* 2006; 13: 491–500.
5. Courneya KS, Segal RJ, McKenzie DC, Dong H, Gelmon K, Friedenreich CM, et al. Effects of exercise during adjuvant chemotherapy on breast cancer outcomes. *Med Sci Sports Exerc* 2014; 46: 1744–1751.
6. Bradshaw PT, Ibrahim JG, Khankari N, Cleveland RJ, Abrahamson PE, Stevens J, et al. Post-diagnosis physical activity and survival after breast cancer diagnosis: the Long Island Breast Cancer Study. *Breast Cancer Res Treat* 2014; 145: 735–742.
7. Hayes SC, Rye S, Disipio T, Yates P, Bashford J, Pyke C, et al. Exercise for health: a randomized, controlled trial evaluating the impact of a pragmatic, translational exercise intervention on the quality of life, function and treatment-related side effects following breast cancer. *Breast Cancer Res Treat* 2013; 137: 175–186.
8. Baumann FT, Bloch W, Weissen A, Brockhaus M, Beulertz J, Zimmer P, et al. Physical activity in breast cancer patients during medical treatment and in the aftercare – a review. *Breast Care* 2013; 8: 330–334.
9. Chlebowski RT. Nutrition and physical activity influence on breast cancer incidence and outcome. *Breast* 2013; 22 Suppl 2: 30–37.
10. Ibrahim EM, Al-Homaidh A. Physical activity and survival after breast cancer diagnosis: meta-analysis of published studies. *Med Oncol* 2011; 28: 753–765.
11. Schmitz KH, Ahmed RL, Hannan PJ, Yee D. Safety and efficacy of weight training in recent breast cancer survivors to alter body composition, insulin, and insulin-like growth factor axis proteins. *Cancer Epidemiol Biomarkers Prev* 2005; 14: 1672–1680.
12. Hayes SC, Johansson K, Stout NL, Prosnitz R, Armer JM, Gabram S, et al. Upper-body morbidity after breast cancer. *Cancer* 2012; 118 (S8): 2237–2249.
13. Courneya KS, McKenzie DC, Mackey JR, Gelmon K, Friedenreich CM, Yasui Y, et al. Effects of exercise dose and type during breast cancer chemotherapy: multicenter randomized trial. *J Natl Cancer Inst* 2013; 105: 1821–1832.
14. Van Vulpen JK, Peeters PH, Velthuis MJ, van der Wall E, May AM. Effects of physical exercise during adjuvant breast cancer treatment on physical and psychosocial dimensions of cancer-related fatigue: a meta-analysis. *Maturitas* 2016; 85: 104–111.
15. Schmidt T, Weisser B, Dürkop J, Jonat W, Van Mackelenbergh M, Röcken C, et al. C. Comparing endurance and resistance training with standard care during chemotherapy for patients with primary breast cancer. *Anticancer Res* 2015; 35: 5623–5629.
16. Do JH, Kim W, Cho YK, Lee J, Song EJ, Chun YM, Jeon JY. Effects of resistance exercise and complex decongestive therapy on arm function and muscular strength in breast cancer related lymphedema. *Lymphology* 2015; 48: 184–196.
17. Cheema BS, Kilbreath SL, Fahey PP, Delaney GP, Atlantis E. Safety and efficacy of progressive resistance training in breast cancer: a systematic review and meta-analysis. *Breast Cancer Res Treat* 2014; 148: 249–268.
18. Sagen Å, Kåresen R, Risberg MA. Physical activity for the affected limb and arm lymphedema after breast cancer surgery. A prospective, randomized controlled trial with two years follow-up. *Acta Oncol* 2009; 48: 1102–1110.
19. Schmitz KH, Troxel AB, Chevillat A, Grant LL, Bryan CJ, Gross CR, et al. Physical activity and lymphedema (the PAL trial): assessing the safety of progressive strength training in breast cancer survivors. *Contemp Clin Trials* 2009; 30: 233–245.
20. Schmitz KH, Ahmed RL, Troxel AB, Chevillat A, Lewis-Grant L, Smith R, et al. Weight lifting for women at risk for breast cancer-related lymphedema. *JAMA* 2010; 304: 2699–2705.
21. Kwan ML, Cohn JC, Armer JM, Stewart BR, Cormier JN. Exercise in patients with lymphedema: a systematic review of the contemporary literature. *J Cancer Surviv* 2011; 5: 320–336.
22. Winters-Stone KM, Lauder milk M, Woo K, Brown JC, Schmitz KH. Influence of weight training on skeletal health of breast cancer survivors with or at risk for breast cancer-related lymphedema. *J Cancer Surviv* 2014; 8: 260–268.
23. Bloomquist K, Karlsmark T, Christensen KB, Adamsen L. Heavy resistance training and lymphedema: prevalence of breast cancer-related lymphedema in participants of an exercise intervention utilizing heavy load resistance training. *Acta Oncol* 2014; 53: 216–225.
24. Ostby P, Armer J, Dale P, van Loo M, Wilbanks C, Stewart B. Surveillance recommendations in reducing risk of and optimally managing breast cancer-related lymphedema. *J Pers Med* 2014; 4: 424–447.
25. Gautam AP, Maiya GA, Vidyasagar MS. Effect of home-based exercise program on lymphedema and quality of life in female postmastectomy patients: pre-post intervention study. *J Rehabil Res Dev* 2011; 48: 1261–1268.
26. Høyer M, Johansson B, Nordin K, Bergkvist L, Ahlgren J, Lidin-Lindqvist A, et al. Health-related quality of life among women with breast cancer – a population-based study. *Acta Oncol* 2011; 50: 1015–1026.
27. Eyigor S, Karapolat H, Yesil H, Uslu R, Durmaz B. Effects of pilates exercises on functional capacity, flexibility, fatigue, depression and quality of life in female breast cancer patients: a randomized controlled study. *Eur J Phys Rehabil Med* 2010; 46: 481–487.
28. Schwarz R, Krauss O, Hinz A. Fatigue in the general population. *Onkologie* 2003; 26: 140–144.
29. Smets EM, Garssen B, Bonke B, De Haes JC. The Multidimensional Fatigue Inventory (MFI) psychometric qualities of an instrument to assess fatigue. *J Psychosom Res* 1995; 39: 315–325.
30. Schmitz KH, Courneya KS, Matthews C, Demark-Wahnefried W, Galvão DA, Pinto BM, et al. American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. *Med Sci Sports Exerc* 2010; 42: 1409–1426.
31. Borg G. Borg's Perceived exertion and pain scales. 1st edn. Human Kinetics. Champaign, 1998.
32. Wahlund HG. Determination of the physical working capacity. A physiological and clinical study with special reference to standardization of cardio-pulmonary functional tests. *Acta Medica Scandinavica Suppl* 215, Stockholm, 1948.
33. Chen X, Zheng Y, Zheng W, Gu K, Chen Z, Lu W, et al. The effect of regular exercise on quality of life among breast cancer survivors. *Am J Epidemiol* 2009; 170: 854–862.
34. Ohira T, Schmitz KH, Ahmed RL, Yee D. Effects of weight training on quality of life in recent breast cancer survivors: the Weight Training for Breast Cancer Survivors (WTBS) study. *Cancer* 2006; 106: 2076–2083.
35. Milne HM, Wallman KE, Gordon S, Courneya KS. Effects of a combined aerobic and resistance exercise program in breast cancer survivors: a randomized controlled trial. *Breast Cancer Res Treat* 2008; 108: 279–288.