



SOCIETAL BURDEN OF STROKE REHABILITATION: COSTS AND HEALTH OUTCOMES AFTER ADMISSION TO STROKE REHABILITATION

Winke VAN MEIJEREN-PONT, MSc^{1,2}, Sietske J. TAMMINGA, PhD^{1,2}, Paulien H. GOOSSENS, MD, PhD³, Iris F. GROENEVELD, PhD⁴, Henk ARWERT, MD, PhD^{1,5}, Jorit J. L. MEESTERS, PhD^{1,2}, Radha RAMBARAN MISHRE, MD^{1,6}, Thea P. M. VLIET VLIELAND, MD, PhD^{1,2} and Wilbert B. VAN DEN HOUT, PhD⁷, on behalf of the Stroke Cohort Outcomes of REhabilitation (SCORE) study group

From the ¹Basalt Rehabilitation, The Hague, ²Department of Orthopaedics, Rehabilitation, and Physical Therapy, Leiden University Medical Center, Leiden, ³Merem Rehabilitation Center, Hilversum, ⁴National Health Care Institute, Diemen, ⁵Department of Rehabilitation Medicine, Haaglanden Medical Centre, The Hague, ⁶Department of Rehabilitation Medicine, Reinier de Graaf Gasthuis, Delft and ⁷Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, The Netherlands

Objective: To estimate societal costs and changes in health-related quality of life in stroke patients, up to one year after start of medical specialist rehabilitation.

Design: Observational.

Patients: Consecutive patients who received medical specialist rehabilitation in the Stroke Cohort Outcomes of REhabilitation (SCORE) study.

Methods: Participants completed questionnaires on health-related quality of life (EuroQol EQ-5D-3L), absenteeism, out-of-pocket costs and healthcare use at start and end of rehabilitation and 6 and 12 months after start. Clinical characteristics and rehabilitation costs were extracted from the medical and financial records, respectively.

Results: From 2014 to 2016 a total of 313 stroke patients completed the study. Mean age was 59 (standard deviation (SD) 12) years, 185 (59%) were male, and 244 (78%) inpatients. Mean costs for inpatient and outpatient rehabilitation were US\$70,601 and US\$27,473, respectively. For inpatients, utility (an expression of quality of life) increased significantly between baseline and 6 months (EQ-5D-3L 0.66–0.73, $p = 0.01$; visual analogue scale 0.77–0.82, $p < 0.001$) and between baseline and 12 months (visual analogue scale 0.77–0.81, $p < 0.001$).

Conclusion: One-year societal costs from after the start of rehabilitation in stroke patients were considerable. Future research should also include costs prior to rehabilitation. For inpatients, health-related quality of life, expressed in terms of utility, improved significantly over time.

Key words: stroke; rehabilitation; cost analysis; utility; health-related quality of life.

Accepted Mar 23, 2021; Epub ahead of print Apr 15, 2021

J Rehabil Med 2021; 53: jrm00201

Correspondence address: Winke van Meijeren-Pont, Basalt Rehabilitation/Department of Orthopaedics, Rehabilitation, and Physical Therapy, Leiden University Medical Center, Wassenaarseweg 501, 2333 AL Leiden, The Netherlands. E-mail: w.pont@basaltrevalidatie.nl

The number of people living with stroke in Europe is expected to increase from 1.1 million per year in 2000 to 1.5 million per year in 2025 (1). Stroke survivors may experience severe functional impair-

LAY ABSTRACT

The objective of this study was to estimate societal costs and changes in health-related quality of life in stroke patients, up to one year after the start of rehabilitation. Participants were stroke patients who received inpatient or outpatient rehabilitation. They completed questionnaires on quality of life, absenteeism, out-of-pocket costs and healthcare use at start and end of rehabilitation and 6 and 12 months after the start of rehabilitation. Rehabilitation costs were obtained from the financial records. From 2014 to 2016 a total of 313 patients completed the study. Mean age was 59 years, 185 (59%) were male and 244 (78%) inpatients. Mean costs for inpatient and outpatient rehabilitation were \$70,601 and \$27,473, respectively. For inpatients, health-related quality of life increased significantly between baseline and 6 months, and between baseline and 12 months. In conclusion, societal costs one year after the start of rehabilitation were considerable and health-related quality of life improved for inpatients.

ments, including impairments in physical functioning (2), cognition (3), and speech/language (4), which, in turn, lead to limitations in activities and participation and to worse quality of life (QoL) (5). Specialist rehabilitation was proven to be effective in improving functional outcomes after stroke (6), such as motor function, balance, walking speed and activities of daily living (7–9). Furthermore, in stroke patients admitted for inpatient rehabilitation, QoL increased significantly between admission and discharge (10).

Besides the fact that rehabilitation after stroke is effective, rehabilitation was also found to be the main contributor to the costs of post-stroke care, according to a systematic review published in 2018 including 42 publications (11). Costs of post-stroke care, but not those of acute care, were included. Rehabilitation in different care settings was evaluated, which included primary, secondary and tertiary care, and the costs often applied to part of the patients and were not described in detail. For the delivery of value-based healthcare (VBHC), it is important to consider not only the health effects and patient-reported outcome measures, but to also evaluate the costs of care, since it is important to achieve good patient outcomes per dollar spent (12, 13).

The aim of the current study was therefore: (i) to estimate the 1-year societal costs from the start of the rehabilitation in stroke patients treated in a medical specialist rehabilitation facility in The Netherlands; and (ii) to evaluate health changes in terms of utility (an expression of quality of life) over that year.

METHODS

Design, setting and subjects

This study was part of the Stroke Cohort Outcomes of REhabilitation (SCORE) study; a longitudinal inception cohort study, which is executed in one secondary care rehabilitation facility with multiple locations in the Netherlands. This study has been described extensively elsewhere (14).

In the Netherlands, after a mean of 8 days of hospital admission, approximately 71% of patients are discharged home, 15% are discharged to geriatric rehabilitation, and 14% are referred to inpatient rehabilitation in a medical specialist rehabilitation facility (15). In general, younger, pre-stroke more active patients with complex impairments are admitted to medical specialist rehabilitation compared with geriatric rehabilitation (16). Furthermore, patients referred to medical specialist rehabilitation want to regain a high level of participation, including return to work, family and social roles and leisure activities. Many different disciplines are involved in medical specialist rehabilitation treatment, and the rehabilitation facilities comprise, amongst others, a sports hall and a swimming pool.

Stroke patients are admitted to the rehabilitation facility for inpatient rehabilitation if they: (i) have had a recent stroke preventing them from living independently at home; (ii) are able to take part in at least 2 therapy sessions of 30 min each per day; (iii) are likely to benefit from rehabilitation therapy; and (iv) are expected to live independently after discharge, whether or not with spouse or caregiver. Stroke patients receive outpatient rehabilitation if they meet the same criteria, but are able to live at home. Stroke patients were eligible for the study, if they were at least 18 years old and had a first or recurrent stroke no longer than 6 months previously. Exclusion criteria were being unable to complete questionnaires in Dutch or not providing written informed consent.

This trial is registered at the Dutch Clinical Trial Registration (NL4147 at www.trialregister.nl). The study protocol of the SCORE-project was approved by the Medical Ethics Board of Leiden University Medical Center (LUMC), P13.249, and is reported in accordance with the STROBE guidelines (17).

Assessments

Patients completed questionnaires at the start of the rehabilitation (baseline), at discharge (inpatients) or at the end of the rehabilitation (outpatients) and at 6 and 12 months after baseline. Appendix 1 shows which questionnaires were completed at the different measurement moments. Clinical characteristics and the Barthel Index (BI) were extracted from the patients' medical file. The BI is a nurse-reported 10-item measurement instrument that scores independence in activities of daily living (ADL) and yields a score between 0 and 20, with higher scores indicating more independence (18).

Of the Stroke Impact Scale (SIS) (19), patients completed the domains communication (7 items), mobility (9 items), memory

and thinking (7 items) and hand functioning (5 items). Items were scored on a 1–5-point Likert scale and transformed to a score out of 100 (19), with higher scores indicating a lower level of difficulty experienced with the task. Internal consistency (Cronbach's α 0.86–0.98) was found to be excellent among stroke survivors and validity was supportive (20).

Healthcare and non-healthcare costs

Societal costs were estimated from the start of the rehabilitation until one year later, separately for inpatients and outpatients. Rehabilitation costs included length of stay in the rehabilitation facility (the number of days for which nursing care was provided) and direct hours of therapy. Volumes and unit prices were obtained from the (financial) administration of the rehabilitation facility. In the patient questionnaires at 6 and 12 months other cost items in the preceding 6 months were assessed. This included healthcare usage outside the rehabilitation facility, out-of-pocket expenses (e.g. for crutches or an electric scooter), informal care, paid home-care and absenteeism. These items were valued using reference prices obtained from the Dutch guidelines for economic evaluations in healthcare (21). If no reference price was available, market prices were used. Absenteeism was valued using the friction costs method, which counts absenteeism for, at most, the duration of the friction period, i.e. the 12-week period considered necessary to fill a vacancy due to long-term sick leave (21). Costs were converted to US dollars (USD) using the purchasing power parity, as listed by the Organisation for Economic Co-operation and Development (OECD) on its website, and are reported at price level 2019 (22). Appendix 1 shows the unit costs.

Health-related quality of life and utility

The patient-reported EuroQol EQ-5D-3L (23) measures health-related quality of life (HRQoL) and consists of 5 domains: mobility, self-care, usual activities, pain/discomfort and anxiety/depression. A visual analogue scale (VAS) records the patient's self-rated health on a vertical scale with endpoints labelled "Best imaginable health state" and "Worst imaginable health state" (23). Utility scores were calculated from the 5 domains using the Dutch tariff (24) and from the VAS scale. A utility or weight of one reflects complete health, whereas 0 reflects health as poor as death (25). The EQ-5D-3L has shown reasonable validity and reliability (23, 26) and moderate responsiveness (27) for patients with stroke.

Statistical analyses

Data analyses were performed in IBM SPSS version 22 v02 (IBM Corp., Armonk, NY, USA, 2013). To account for systematic missingness, data were imputed using multiple imputation by chained equations (MICE) (28, 29) with predictive mean matching (30, 31) and 100 imputation sets. Missing values for out-of-pocket costs were imputed, based on either the mean price if 50% or more of the participants filled in a price, or market prices otherwise.

Characteristics of patients who did and did not agree to participate, who did and did not complete the study period and inpatients and outpatients were compared using independent-sample *t*-tests, Fisher's exact tests or Mann-Whitney *U* tests, where appropriate. Utility scores were compared using paired-sample *t*-tests at baseline vs 6 months, baseline vs 12 months, and 6 vs 12 months, respectively. Univariate linear regression

analyses with total costs as a function of baseline utility were performed separately for inpatients and outpatients.

RESULTS

Fig. 1 shows that, between 10 March 2014 and 31 August 2016, 791 stroke patients, were admitted for inpatient or outpatient rehabilitation. Of these, 182 (23%) patients were missed and 609 (77%) patients were invited to participate. A total of 244 (40%) of the invited patients were not willing to participate. A total of 365 (60%) of the invited patients signed informed consent and completed one or more questionnaires. Sex and age did not differ significantly between patients who did and did not participate in the study (mean age 59.7 vs 60.4 years, $p=0.40$; percentage male 58% vs 56%, $p=0.66$, respectively).

The 12-month follow-up period was completed by 313 patients. Fifty-two patients dropped out (17%). Baseline characteristics did not differ significantly between patients who did and did not complete the study (see Table I).

Of those who completed the study, inpatients were significantly older than outpatients (mean age 60 vs 56 years, $p=0.02$). Furthermore, inpatients scored significantly better than outpatients on communication (median 92 vs 86, $p=0.01$) and memory and thinking (median 89 vs 75, $p<0.01$) as measured with the SIS at baseline. Outpatients had significantly better hand function than inpatients at baseline (median 75 vs 50, $p=0.02$).

A total of 244 (78%) patients received inpatient rehabilitation, with a median duration of 44 days (interquartile range (IQR) 14–155). Of these, 160 (61%) received outpatient rehabilitation thereafter, with a median duration of 126 days (IQR 72–186). The median duration of rehabilitation for outpatients was 105 days (IQR 70–164).

Of the baseline measurements 5% were missing. Of the QoL, absenteeism, and healthcare use measurements at baseline 12%, 38% (in patients younger than 66 years, who reported that they had paid work at baseline) and 2% were missing.

Costs

Mean total costs were US\$70,601 for inpatients and \$27,473 for outpatients (see Table II). For inpatients, rehabilitation was the biggest contributor to costs (\$46,870; 66%), followed by productivity loss (\$10,211; 14%) and informal care (\$6,575; 9%). For outpatients, rehabilitation was also the biggest contributor (\$9,899; 36%), although to a lesser extent than in inpatients, followed by productivity loss (\$9,416; 34%) and informal care (\$4,531; 16%).

The costs of rehabilitation therapy for inpatients were approximately 3 times higher than for outpatients (\$30,741 vs \$9,899). The largest contributor to the costs of rehabilitation for outpatients was physical therapy (\$2,579; 26% of the total rehabilitation costs), followed by occupational therapy (\$2,061; 21%) and psychology (\$1,835; 19%). For inpatients the costs of stay were the largest contributor to the costs of rehabilitation (\$16,129; 34%), followed by physical therapy (\$7,707; 16%) and occupational therapy (\$5,813; 12%).




Fig. 1. Flowchart of stroke patients included in the Stroke Cohort Outcomes of REhabilitation (SCORE) study between March 2014 and August 2016.

Table I. Characteristics of stroke patients admitted for inpatient or outpatient rehabilitation in a rehabilitation facility, completing and not completing the present study

	Inpatients who completed the study, n = 244	Outpatients who completed the study, n = 69	p-value ^a inpatients vs outpatients	Patients who completed the study, n = 313	Patients who did not complete the study, n = 52	p-value ^c patients who did vs who did not complete the study
Sex, male, n (%)	146 (60)	39 (57)	0.62	185 (59)	28 (54)	0.54
Age, years, mean (SD)	60 (12)	56 (12)	0.02	59 (12)	62 (12)	0.11
Education level, n (%)			0.19			0.41
Low	109 (45)	22 (32)		131 (42)	26 (50)	
Medium	63 (26)	26 (38)		88 (28)	13 (25)	
High	72 (30)	22 (32)		94 (30)	14 (27)	
Type of stroke (ischaemic), n (%)	183 (75)	57 (83)	0.20	240 (77)	41 (79)	0.71
Stroke localization, n (%)			0.37			0.66
Left	109 (45)	38 (57)		147 (48)	26 (50)	
Right	104 (43)	22 (33)		127 (41)	21 (40)	
Other	31 (13)	9 (12)		40 (13)	5 (9)	
Rehabilitation (inpatient), n (%)				244 (78)	45 (87)	0.20
Duration inpatient rehabilitation, days, median (IQR)	44 (14–155) n = 239	–				
Outpatient rehabilitation after inpatient rehabilitation, n (%)	146 (61) n = 241	–				
Duration outpatient rehabilitation, days, median (IQR)	126 (72–186) n = 113	105 (70–164) n = 67				
Barthel Index, mean (SD) ^a	14 (5)	–	–	14 (4)	14 (5)	0.54
Communication (SIS) ^b , median (IQR)	93 (79–100) n = 212	86 (71–93) n = 64	0.01	92 (75–100) n = 282	88 (73–96) n = 45	0.43
Mobility (SIS) ^b , median (IQR)	81 (43–96) n = 37	92 (78–100) n = 20	0.06	83 (64–97) n = 57	58 (54–92) n = 7	0.15
Memory and thinking (SIS) ^b , median (IQR)	89 (74–96) n = 218	75 (65–86) n = 64	<0.01	86 (71–96) n = 288	80 (63–94) n = 46	0.24
Hand function (SIS) ^{b,d} , median (IQR)	50 (5–80) n = 130	75 (55–80) n = 20	0.02	60 (9–80) n = 150	40 (3–73) n = 24	0.30

^aScale range 1–20; higher scores denote better functioning. ^bScale range 0–100; higher scores indicate a lower level of difficulty experienced with the task. ^cp-value of the independent samples t-test, the Fisher's exact test or the Mann-Whitney U test, where appropriate. ^dOnly for patients who indicated that their hand was affected by stroke. SD: standard deviation.

Table II. Mean resource use and costs among stroke inpatients and outpatients, in the first year after admission to a rehabilitation facility

	Inpatients, n = 244			Outpatients, n = 69		
	% patients	Volume ^a	Costs (US\$) ^b	% patients	Volume ^a	Costs (US\$) ^b
Rehabilitation						
Rehabilitation physician, h	100	10.7	4,078	100	1.9	833
Physical therapy, h	100	47.2	7,707	96	16.2	2,579
Occupational therapy, h	100	36.4	5,813	99	13.2	2,061
Speech-language therapy, h	100	18.4	3,014	70	9.2	1,073
Psychology, h	99	16.3	4,082	86	14.6	1,835
Social work, h	100	12.3	2,010	96	8.2	1,271
Recreational therapy, h	98	8.2	1,155	4	2.1	11
Other therapy, h	100	23.5	2,881	51	2.4	236
Total rehabilitation therapy, mean (SD)			30,741 (17,345)			9,899 (6,020)
Rehabilitation stay, days	100	50.0	16,129	–	–	–
Total rehabilitation, mean (SD)			46,870 (24,580)			9,899 (6,020)
Non-rehabilitation healthcare						
Hospital readmissions, days	22	15.5	2,046	16	8.2	785
General practitioner, visits	81	5.5	185	79	5.4	179
Neurologist, visits	56	2.3	158	65	2.3	186
Other medical specialists, visits	63	3.9	277	68	6.5	505
Occupational physician, visits	33	4.5	252	43	5.3	384
Allied health professionals, visits	67	32.9	1,082	52	22.0	566
Total non-rehabilitation healthcare, mean (SD)			3,999 (7,000)			2,604 (2,940)
Total healthcare costs, mean (SD)			50,869 (26,617)			12,502 (7,023)
Other non-healthcare						
Out of pocket costs, number of devices	65	4.3	2,575	28	2.4	889
Informal care, h	80	469.1	6,575	69	374.1	4,531
Paid home care, h	11	88.8	371	4	54.7	135
Productivity loss, h ^c	42	375.2	10,211	60	271.6	9,416
Total non-healthcare costs, mean (SD)			19,732 (15,267)			14,970 (12,244)
Total costs, mean (SD)			70,601 (34,534)			27,473 (15,200)

^aVolume for patients who received care. ^bCosts for total population. ^cProductivity loss was calculated for patients under the age of 65 years who reported that they had paid work at baseline. SD: standard deviation.




Fig. 2. Utility scores calculated from the EuroQoL EQ-5D-3L classification system and from the visual analogue scale (VAS), at start of rehabilitation, 6 and 12 months for stroke patients admitted to a rehabilitation facility. *Statistically significant differences compared with start of rehabilitation.

Utility scores

Fig. 2 shows the utility scores over time, according to the EQ-5D-3L and the VAS. For inpatients, mean baseline utility was 0.66 (standard deviation (SD) 0.27) from the EQ-5D-3L and 0.77 (SD 0.16) from the VAS. For outpatients, mean baseline utility was 0.69 (SD 0.23) from the EQ-5D-3L and 0.79 (SD 0.15) from the VAS. For inpatients, utility improved significantly between baseline and 6 months (EQ-5D-3L, $p=0.01$; VAS $p<0.001$) and between baseline and 12 months (VAS, $p<0.001$). For outpatients there was no statistically significant change over time. The decrease in utility according to the EQ-5D-3L between 6 and 12 months observed in inpatients was not statistically significant ($p=0.11$).

Tables III and IV show the results of the linear regression analyses with total costs as a dependent variable and utility as an independent variable. Baseline utility from the EQ-5D-3L and VAS are both significantly associated with total costs for inpatients ($p<0.001$ for both). For outpatients baseline utility from the VAS was significantly associated with total costs ($p=0.014$). For example, an outpatient with a

Table III. Linear regression analysis of total costs as the dependent variable and utilities as independent variable, for stroke patients who received inpatient rehabilitation ($n = 244$) in a rehabilitation facility

Inpatient model	Point estimate	95% CI	p-value
Baseline EQ-5D-3L			
Intercept	98,091	85,227 to 110,954	<0.001
Slope	-41,526	-59,369 to -23,682	<0.001
Baseline VAS			
Intercept	144,426	122,723 to 166,128	<0.001
Slope	-96,010	-123,616 to -68,404	<0.001

CI: confidence interval; VAS: visual analogue scale; EQ-5D-3L: EuroQoL EQ-5D-3L.

Table IV. Linear regression analysis of total costs as the dependent variable and utilities as independent variable, for stroke patients who received outpatient rehabilitation ($n = 69$) in a rehabilitation facility

Outpatient model	Point Estimate	95% CI	p-value
Baseline EQ-5D-3L			
Intercept	29,844	17,725 to 41,964	<0.001
Slope	-3,408	-20,020 to 13,205	0.688
Baseline VAS			
Intercept	58,162	38,755 to 77,568	<0.001
Slope	-38,934	-63,115 to -14,753	0.002

95% CI: 95% confidence interval.

baseline VAS utility score of 0.79 is expected to have total costs of $58,162 - 38,934 \times 0.79 = 27,404$ USD, whereas an outpatient with a worse baseline VAS utility score of 0.49 has higher expected total costs, of $58,162 - 38,934 \times 0.49 = 39,084$ USD.

DISCUSSION

One-year costs after the start of medical specialist rehabilitation post stroke from a societal perspective were \$70,601 and \$27,473 for inpatients and outpatients, respectively. For both inpatients and outpatients, rehabilitation was the biggest contributor, yet to a larger extent in inpatients than in outpatients. Both the costs for stay in the rehabilitation facility and for all types of therapy were higher. Productivity loss and informal care were other large contributors to the costs for both inpatients and outpatients. Between baseline and 6 months, and baseline and 12 months, utility improved significantly for inpatients. A linear regression analysis showed that utility at baseline significantly predicted costs.

Communication, memory and thinking and hand function differed significantly between inpatients and outpatients in the current study. Patients with motor problems were more often admitted for inpatient rehabilitation, whereas patients with cognitive complaints more often received outpatient rehabilitation. Using the EuroQoL EQ-5D-3L and VAS, significant improvements were found over time in inpatients, but not in outpatients. This could be explained by the smaller number of outpatients or, alternatively, by the fact that the EQ-5D-3L does not explicitly measure cognitive complaints, which are more prevalent among outpatients. The differences between inpatients and outpatients might also partly explain the higher costs of rehabilitation treatment for inpatients. Since the clinical characteristics of inpatients and outpatients differ significantly at admission, it is not valid to compare outcomes in terms of utilities and costs between these groups.

Comparison with the literature

The current results are in line with a previous review, which found that rehabilitation was the main contri-

butor to the costs of post-stroke care, followed by informal care (11). In other studies the costs of medical interventions, physiotherapy, occupational and speech therapy, nursing care, primary care visits, readmissions to hospital due to recurrent stroke, emergency care during the rehabilitation period and other costs, such as medication, community services, transportation, meals on wheels and assistive devices, were included. Although the costs for rehabilitation found in the present study are high, research showed that the benefits for society outweigh the costs (32).

At 6 and 12 months after the start of the rehabilitation EQ-5D-3L utility scores were 0.73 and 0.69 for inpatients and 0.74 and 0.74 for outpatients. These results are mostly in line with a Dutch hospital-based study that found utility scores of 0.74 and 0.74 at 6 and 12 months post stroke, respectively (15). The lower 0.69 utility score for inpatients at 12 months in the current study might be explained by the fact that patients referred to medical specialist rehabilitation are more severely affected by stroke than patients included in a hospital-based study (16).

Rehabilitation facilities in the Netherlands are obliged to work with national guidelines. Yet the recommendations are not very detailed, leaving room for local variation. Previous research showed that there were many similarities, but to some extent there was also some practice variation in the structure and processes of rehabilitation, as delivered by rehabilitation facilities (33, 34). Variation mainly concerned patient subgroups, clinical pathways and the duration of aftercare (34). Practice variation might lead to some difference in costs. However, differences are expected to be small, since health insurers and healthcare providers have made agreements on the price of healthcare, based on the amount of care a patient with a certain diagnosis needs on average. This is also the case for stroke rehabilitation.

Strengths and limitation

A strong point of this study is that there are not many studies on costs of medical specialist rehabilitation that include patient-reported out-of-pocket costs, absenteeism, healthcare usage and utility. Furthermore, different types of therapy during rehabilitation were estimated separately. Evaluating the costs of care is an important aspect in the delivery of value-based healthcare (12, 13).

An important limitation of the present study is that the included costs started at the start of the rehabilitation. Already before the start of the rehabilitation considerable costs are incurred, for example for ambulance care, emergency care, hospital stay, magnetic reso-

nance imaging (MRI), thrombolysis or thrombectomy. Costs in the Netherlands may also not be representative for other healthcare setting. Patient reports about absenteeism and the EQ-5D-3L contained more than 10% missing values. Multiple imputation was used to account for bias, but may not have prevented all bias.

An additional limitation was that absenteeism was self-reported, possibly leading to under-reporting of the time someone was absent (35). Of all patients eligible for participation in this study, 54% were missed or not willing to participate. Therefore, selection bias may have occurred. Although sex and age did not differ significantly between patients who participated in the study and those who did not because they were not invited or refused participation, a limitation of this study is that we do not know whether these patients differed on other characteristics, such as functional limitations. Patients who have more functional or cognitive limitations might not have been able to participate in the study and might need more time to learn and therefore more care. The results of the regression analysis showed that patients with a worse baseline utility have higher total expected costs. Given this lack of information, it remains unclear whether the costs found in the present study could be somewhat over- or under-estimated.

Recommendations for future research

Costs prior to admission to the rehabilitation facility and costs of medication were not included in the present study. For future research it would be recommended to include these costs.

The European Stroke Organisation (ESO) Health Economics Working Group made a protocol to standardize and improve the economic evaluations of interventions for stroke. Resources mentioned in this protocol, but not included in the present study, were amongst others transport, change in residence and living arrangements, medications and more clinical outcomes after treatment (36). Although this was not a comparative study, gathering this data on resources might help to standardize research and compare outcomes. Therefore, in future research it would be recommended to gather data on these resources.

Another recommendation for future research is to consider extracting healthcare usage outside the rehabilitation facility from a health insurer or other central administration system in order to get more complete data. Such administrative data do not rely on patients' recall, but can be difficult to obtain, and may lack the detail necessary to provide real insight (37). Previous research on the reliability of stroke patients' reports of general practitioner visits over 12 months found

that patients modestly under-reported the number of visits (38).

CONCLUSION

In conclusion 1-year costs from the start of medical specialist rehabilitation post stroke from a societal perspective were estimated at US\$70,601 and \$27,473 for inpatients and outpatients, respectively. Future research should include costs prior to the rehabilitation, since considerable costs are incurred in the acute phase. For inpatients, utility improved significantly between the start of the rehabilitation and both 6 and 12 months.

Funding. This work was supported by the Stichting Kwaliteitsgeld Medisch Specialisten.

The authors have no conflicts of interest to declare.

REFERENCES

1. Truelsen T, Piechowski-Jóźwiak B, Bonita R, Mathers C, Bogousslavsky J, Boysen G. Stroke incidence and prevalence in Europe: a review of available data. *Eur J Neurol* 2006; 13: 581–598.
2. Langhorne P, Coupar F, Pollock A. Motor recovery after stroke: a systematic review. *Lancet Neurol* 2009; 8: 741–754.
3. van Rijsbergen MW, Mark RE, de Kort PL, Sitskoorn MM. Subjective cognitive complaints after stroke: a systematic review. *J Stroke Cerebrovasc Dis* 2014; 23: 408–420.
4. Lazar RM, Boehme AK. Aphasia as a predictor of stroke outcome. *Curr Neurol Neurosci Rep* 2017; 17: 83.
5. Carod-Artal J, Egido JA, González JL, Varela de Seijas E. Quality of life among stroke survivors evaluated 1 year after stroke: experience of a stroke unit. *Stroke* 2000; 31: 2995–3000.
6. Ottenbacher KJ, Jannell S. The results of clinical trials in stroke rehabilitation research. *Arch Neurol* 1993; 50: 37–44.
7. Veerbeek JM, van Wegen E, van Peppen R, van der Wees PJ, Hendriks E, Rietberg M, et al. What is the evidence for physical therapy poststroke? A systematic review and meta-analysis. *PLoS One* 2014; 9: e87987.
8. Pollock A, Baer G, Campbell P, Choo PL, Forster A, Morris J, et al. Physical rehabilitation approaches for the recovery of function and mobility following stroke. *Cochrane Database Syst Rev* 2014; 4: CD001920.
9. Legg LA, Lewis SR, Schofield-Robinson OJ, Drummond A, Langhorne P. Occupational therapy for adults with problems in activities of daily living after stroke. *Cochrane Database Syst Rev* 2017; 7: CD003585.
10. Groeneveld IF, Goossens PH, van Meijeren-Pont W, Arwert HJ, Meesters JJJ, Rambaran Mishre AD, et al. Value-based stroke rehabilitation: feasibility and results of patient-reported outcome measures in the first year after stroke. *J Stroke Cerebrovasc Dis* 2019; 28: 499–512.
11. Rajsic S, Gothe H, Borba HH, Sroczynski G, Vujicic J, Toell T, et al. Economic burden of stroke: a systematic review on post-stroke care. *Eur J Health Econ* 2019; 20: 107–134.
12. Porter ME. What is value in health care? *N Engl J Med* 2010; 363: 2477–2481.
13. Porter ME, Olmsted Teisberg E. Redefining health care. Creating value-based competition on results. Boston: Harvard Review Business Press; 2006.
14. Groeneveld IF, Meesters JJJ, Arwert HJ, Rambaran Mishre AD, Vliet Vlieland TPM, Goossens PH. Onderzoeksopzet met analyse van structuur, proces en uitkomsten: Praktijkvariatie in de CVA-revalidatie. [Research design of an analysis of structure, processes and outcomes: Practice variation in stroke rehabilitation.] *Nederlands Tijdschrift voor Revalidatiegeneeskunde* 2015; 3: 134–137 (in Dutch).
15. van Eeden M, van Heugten C, van Mastrigt GA, van Mierlo M, Visser-Meily JM, Evers SM. The burden of stroke in the Netherlands: estimating quality of life and costs for 1 year poststroke. *BMJ Open* 2015; 5: e008220.
16. de Boer AR, van Dis I, Visseren FLJ, Vaartjes I, Bots ML. Hart- en vaatziekten in Nederland, cijfers over leefstijl- en risicofactoren, ziekte en sterfte [Cardiovascular diseases in the Netherlands, numbers on lifestyle and risk factors, illness and mortality.] Den Haag: Nederlandse Hartstichting; 2018 (in Dutch).
17. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies. *Int J Surg* 2014; 12: 1495–1499.
18. Quinn TJ, Langhorne P, Stott DJ. Barthel Index for stroke trials: development, properties, and application. *Stroke* 2011; 42: 1146–1151.
19. Duncan PW, Wallace D, Lai SM, Johnson D, Embretson S, Laster LJ. The stroke impact scale version 2.0. Evaluation of reliability, validity, and sensitivity to change. *Stroke* 1999; 30: 2131–2140.
20. Vellone E, Savini S, Fida R, Dickson VV, Melkus GD, Carod-Artal FJ, et al. Psychometric evaluation of the Stroke Impact Scale 3.0. *J Cardiovasc Nurs* 2015; 30: 229–241.
21. IJzerman MJ, Al MJ, de Boer A, Brouwer WBF, van Busschbach JJ, Dijkgraaf MGW, et al. 2015. Richtlijn voor het uitvoeren van economische evaluaties in de gezondheidszorg. [Costing manual: methodology of costing research and reference prices for economic evaluations in healthcare.] Diemen: Zorginstituut Nederland; 2015 (in Dutch).
22. OECD. Purchasing power parities (PPP) (indicator). 2019[cited 2019 Oct 24]. Available from: https://www.oecd-ilibrary.org/finance-and-investment/purchasing-power-parities-ppp/indicator/english_1290ee5a-en
23. Lamers LM, McDonnell J, Stalmeier PF, Krabbe PF, Busschbach JJ. The Dutch tariff: results and arguments for an effective design for national EQ-5D valuation studies. *Health Econ* 2006; 15: 1121–1132.
24. Drummond MF, O'Brien B, Stoddart GL, Torrance GW. Methods for the economic evaluation of health care programmes. New York: Oxford University Press; 1997.
25. Hunger M, Sabariego C, Stollenwerk B, Cieza A, Leidl R. Validity, reliability and responsiveness of the EQ-5D in German stroke patients undergoing rehabilitation. *Qual Life Res* 2012; 21: 1205–1216.
26. Lu WS, Huang SL, Yang JF, Chen MH, Hsieh CL, Chou CY. Convergent validity and responsiveness of the EQ-5D utility weights for stroke survivors. *J Rehabil Med* 2016; 48: 346–351.
27. Golicki D, Niewada M, Karlińska A, Buczek J, Kobayashi A, Janssen MF, et al. Comparing responsiveness of the EQ-5D-5L, EQ-5D-3L and EQ VAS in stroke patients. *Qual Life Res* 2015; 24: 1555–1563.
28. van Buuren S, Oudshoorn K. Flexible multiple imputation by MICE. Leiden: TNO Prevention and Health; 1999.
29. van Buuren S. Multiple imputation of discrete and continuous data by fully conditional specification. *Stat Methods Med Res* 2007; 16: 219–242.
30. Rubin DB. Multiple imputation for nonresponse in surveys. New York: Wiley; 1987.
31. Little RJ, Yosef M, Cain KC, Nan B, Harlow SD. A hot-deck multiple imputation procedure for gaps in longitudinal data on recurrent events. *Stat Med* 2008; 27: 103–120.
32. Kok L, Houkes A, Niessen N. Kosten en baten van revalidatie [Costs and benefits of rehabilitation.] Amsterdam: SEO Economisch Onderzoek; 2008 (in Dutch).

33. Goossens PH, Groeneveld IF, van Vree FM, Arwert H, Ramban-Mishre AD, Meesters JLL, et al. Uitkomsten van het Stroke Cohort Outcome of REhabilitation (SCORE)-onderzoek. Praktijkvariatie in de CVA-revalidatie. [Outcomes of the Stroke Cohort Outcome of REhabilitation (SCORE)-study. Practice variation in stroke rehabilitation.] *Nederlands Tijdschrift voor Revalidatiegeneeskunde* 2018; 3: 124–128 (in Dutch).

34. Groeneveld IF, Meesters JJ, Arwert HJ, Roux-Otter N, Ribbers GM, van Bennekom CA, et al. Practice variation in the structure of stroke rehabilitation in four rehabilitation centres in the Netherlands. *J Rehabil Med* 2016; 48: 287–292.

35. Johns G, Miraglia M. The reliability, validity, and accuracy of self-reported absenteeism from work: a meta-analysis. *J Occup Health Psychol* 2015; 20: 1–14.

36. Cadilhac DA, Kim J, Wilson A, Berge E, Patel A, Ali M, et al. Improving economic evaluations in stroke: a report from the ESO Health Economics Working Group. *Eur Stroke J* 2020; 5: 184–192.

37. van den Brink M, van den Hout WB, Stiggelbout AM, van de Velde CJ, Kievit J. Cost measurement in economic evaluations of health care: whom to ask? *Med Care* 2004; 42: 740–746.

38. Chishti T, Harris T, Conroy R, Oakeshott P, Tulloch J, Coster D, et al. How reliable are stroke patients' reports of their numbers of general practice consultations over 12 months? *Fam Pract* 2013; 30: 119–122.

Appendix 1. Questionnaires completed by stroke patients and other data sources, measurement moments and unit costs

	Source	Start of rehabilitation	Discharge/end of rehabilitation	6 months	12 months	Unit costs (US\$)
<i>Sociodemographic characteristics</i>						
Sex	Patient reported	x				
Age						
Education level (low, medium, high)						
<i>Clinical characteristics</i>						
Type of stroke (ischaemic)	Medical file	x	x			
Stroke localization (left, right, other)						
Rehabilitation (inpatient)						
Duration inpatient rehabilitation						
Outpatient rehabilitation after inpatient rehabilitation						
Duration outpatient rehabilitation						
Barthel Index						
Stroke Impact Scale (SIS)	Patient reported	x				
Health-related quality of life and utility						
EQ-5D-3L and visual analogue scale	Patient reported	x		x	x	
Healthcare usage and costs						
Direct h of therapy rehabilitation facility (rehabilitation physician, physical therapist, sports therapist, occupational therapy, speech-language therapist, clinical linguist, psychologist, psychology assistant, psychiatrist, social work, recreational therapy, other therapy)	Administration system				x	Range 85–402
Length of stay rehabilitation facility	Administration system				x	326
Hospital readmissions	Patient reported			x	x	597
General practitioner						41
Neurologist						124
Other medical specialists						114
Occupational physician						169
Allied health professionals (physical therapist, occupational therapist, speech and language therapist, psychologist, social worker, dietician, sexologist)						Range 38–112
<i>Non-healthcare costs</i>						
Out of pockets costs	Patient reported		x	x	x	
Informal care	Patient reported			x	x	18
Paid home care	Patient reported			x	x	
Domestic help						25
Care						63
Nursing						92
Productivity loss	Patient reported	x		x	x	44